- tok to tok tok	414 Rec'd	PCT/PTO 20 SED BOOK T \$
	MERCE PATER TAND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER
(REV 12-29-99) TRANSMITTAL LETTER	TO THE UNITED STATES	0623.0890000/EKS/PSC
DESIGNATED/ELECTE		U.S. APPLICATION NO (IF KNOWN, SEE 37 C F.R § 1 5)
CONCERNING A FILING	` '	To be a 1994 646679
INTERNATIONAL APPLICATION NO	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/GB99/00905	22 March 1999	20 March 1998
TITLE OF INVENTION		
Signal Transduction Protein Involved in Pl	ant Dehiscence	
APPLICANT(S) FOR DO/EO/US Wyatt, Paul; Roberts, Jeremy Alan; and Wi	hitelaw Catherine	
		US) the following items and other information:
Applicant herewith sublints to the Officer S	tates Designated Diceted Office (DO/DO/	obj the following fields and other information.
1. 🛮 This is a FIRST submission of ite	ems concerning a filing under 35 U.S.C. 3	71.
2. This is a SECOND or SUBSEQU	JENT submission of items concerning a fi	ling under 35 U.S.C. 371.
		371(f)) at any time rather than delay examination until the
	limit set in 35 U.S.C. 371(b) and PCT Art	
· ·	•	the 19th month from the earliest claimed priority date.
<u></u>	ication as filed (35 U.S.C. 371(c)(2))	
· · · · · · · · · · · · · · · · · · ·	(required only if not transmitted by the In	ternational Bureau).
	the International Bureau.	
c. Li s not required, as the a	pplication was filed in the United States R	Leceiving Office (RO/US).
A translation of the International	Application into English (35 U.S.C. 371(c)(2)).
7. 🛛 Amendments to the claims of the	International application under PCT Artic	cle 19 (35 U.S.C. 371(c)(3))
** <u>*</u>	h (required only if not transmitted by the I	nternational Bureau).
b. have been transmitted by	by the International Bureau.	
c. have not been made; he	owever, the time limit for making such am	endments has NOT expired.
d. Mave not been made an	d will not be made.	
	to the claims under PCT Article 19 (35 U.	S.C. 372(c)(3)).
9. An oath or declaration of the inve	entor(s) (35 U.S.C. 371(c)(4)).	
All	ninary Examination Report under PCT Ar	ticle 36 (35 U.S.C. 371(c)(5)).
Items 11. to 16. below concern other doc	ument(s) or information included:	
11. An Information Disclosure States	ment under 37 C.F.R. 1.97 and 1.98.	
12. An assignment document for reco	ording. A separate cover sheet in complia	nce with 37 C.F.R. 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendmen	t and submission of sequence listing.	
☑ A SECOND or SUBSEQUENT	preliminary amendment.	

16. Other items or information: Authorization To Treat A Reply As Incorporating An Extension Of Time Under 37 C.F.R. § 1.136(a)(3) (in duplicate); and Claim for Priority Under 35 U.S.C. § 119(a)-(d) In Utility Application, with Certified Priority Document attached.

14. A substitute specification.

15. A change of power of attorney and/or address letter.

534 Rec'd PCT/PTC 20 SEP 2000

INTERNATIONAL APPLICATION NO U.S. APPLICATION NO. (if known, see 37 C.F.R. 0623.0890000/EKS/PSC PCT/GB99/00905 1.50) **CALCULATIONS** PTO USE ONLY 17. X The following fees are submitted: Basic National Fee (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$970.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$840.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International preliminary examination fee paid to USPTO (37 CFR 1.482) International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) \$ 96.00 ENTER APPROPRIATE BASIC FEE AMOUNT \$840.00 \$ 130.00 Surcharge of \$130.00 for furnishing the oath or declaration later than \Box 20 ⊠ 30 months from the earliest claimed priority date (37 CFR 1.492(e)). Claims Number Filed Number Extra Rate Fotal Claims 29 - 20X \$18.00 \$ 162.00 Independent Claims 0 X \$78.00 \$ 0.00 + \$260.00 \$ 0.00 Multiple dependent claim(s) (if applicable) j TOTAL OF ABOVE CALCULATIONS \$1,132.00 Reduction of ½ for filing by small entity, if applicable. A Small Entity Statement must be 黃led. (Note 37 CFR 1.9, 1.27, 1.28) SUBTOTAL = \$1,132.00 \$ Processing fee of \$130.00 for furnishing the English translation later than \square 20 \square 30 months from the earliest claimed priority date (37 CFR 1.492(f)). TOTAL NATIONAL FEE \$ 1,132.00 Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property TOTAL FEES ENCLOSED = \$1,132.00 Amount to be: \$ refunded: \$ charged: a. 🛛 A check in the amount of \$1,132.00 to cover the above fees is enclosed. in the amount of \$_____ to cover the above fees. A duplicate copy of this Please charge my Deposit Account No. sheet is enclosed. The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit c. X Account No. 19-0036. A duplicate copy of this sheet is enclosed. NOTE: Where an appropriate time limit Under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status. SEND ALL CORRESPONDENCE TO: STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C. Eric K. Steffe 1100 New York Avenue, NW, Suite 600

36,688

Washington, D.C. 20005-3934

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE 2000

In re application of:

Wyatt et al.

Art Unit:

To be assigned

Appl. No. To be assigned (U.S. National

Phase of PCT/GB99/00905)

Examiner:

To be assigned

Filed: Herewith

Atty. Docket: 0623.0890000/EKS/PSC

For: **Signal Transduction Protein**

Involved in Plant Dehiscence

Second Preliminary Amendment

Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

In advance of prosecution please amend the application as follows:

In the Claims

Please amend the claims as follows:

- 3. Nucleic acid as claimed in claim 1 [or claim 2] which is naturally expressed in a dehiscence zone.
 - 4. Nucleic acid as claimed in claim 1 encoding a protein wherein the protein:
 - a) comprises the amino acid sequence shown in Figure 1 or;
 - has one or more amino acid deletions, insertions or substitutions relative b) to a protein as defined in a) above, and has at least 40% amino acid sequence identity therewith; or

- a fragment of a protein as defined in a) or b) above which is at least 10 amino acids long.
- 5. Nucleic acid as claimed in [any one of claims 1 to 4] <u>claim 1</u> which comprises the sequence set out in Figure 1 or a fragment thereof which is at least 30 bases long.
- 6. Nucleic acid, as claimed in [any one of claims 1 to 5] <u>claim 1</u> in combination with one or more further nucleic acid sequence which is dehiscence-zone expressed.
- 7. Nucleic acid which is antisense to nucleic acid as claimed in [any one of claims 1 to 6] claim 1.
- 8. Nucleic acid as claimed in [any one of claims 1 to 7] <u>claim 1</u> including a promoter or other regulatory sequence which controls expression of the nucleic acid.
- 9. Nucleic acid which is the naturally occurring promoter or other regulatory sequence which controls expression of nucleic acid as claimed in [any one claims 1 to 8] <u>claim</u>
 1.
- 10. Nucleic acid as claimed in [any one of claims 1 to 9] <u>claim 1</u> which is in the form of a vector.
 - 11. A cell comprising nucleic acid as claimed in [any one of claims 1 to 10] claim 1.

- 13. A process for obtaining a cell [as claimed in claim 11 or claim 12] comprising introducing nucleic acid as claimed in [any one of claims 1 to 10] claim 1 into said cell.
 - 14. A plant or a part thereof comprising a cell as claimed in [claim 11 or] claim 12.
- 15. Propagating material or a seed comprising a cell as claimed in [claim 11 or] claim 12.
- 16. A process for obtaining a plant or plant part [as claimed in claim 14 or claim 15] comprising obtaining a cell as claimed in claim 11 and growth thereof [or obtaining a plant, plant part, or propagating material as claimed in claim 14 or claim 15 and growth thereof].
 - 18. A protein as claimed in claim 17 which:
 - a) comprises the amino acid sequence shown in Figure 1 or;
 - b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above, and has at least 40% amino acid sequence identity therewith; or
 - a fragment of a protein as defined in a) or b) above which is at least 10 amino acids long.
 - 19. A plant as claimed in claim 17 [or claim 18] which is isolated or recombinant.
- 21. A process [as claimed in claim 20] for regulating or controlling dehiscence in a plant or plant part which comprises obtaining a plant cell as claimed in [claim 21 or part of a plant as claimed in claim 14] claim 12 and deriving a plant therefrom.

- 22. A process <u>for regulating or controlling dehiscence in a plant or plant part</u> [as claimed in claim 20] which comprises obtaining propagating material or a seed as claimed in claim 15 and deriving a plant therefrom.
- 24. A process for controlling or regulating plant dehiscence comprising introducing the [Use of] nucleic acid as claimed in [any one of claims 1 to 10] claim 1 [in the control/regulation of plant dehiscence] into a cell, tissue, plant part thereof or propagating material and expressing said nucleic acid.
- 25. [Use of nucleic] <u>Nucleic</u> acid as claimed in [any one of claims 1 to 10 as] <u>claim</u> 4 which is a probe.
- 26. A process for producing a cell, tissue, plant part thereof or propagating material comprising introducing the [Use of] nucleic acid as claimed in [any one claims 1 to 10] claim 1 into a cell, tissue, plant part thereof or propagating material and causing growth of said cell, tissue, plant part thereof or propagating material [in the production of a cell, tissue, plant part thereof or propagating material].
- 29. [Use of a] A protein as claimed in [any one of claims 17 to 19] claim 18 which is [as] a probe.

Wyatt et al. Appl. No. To be assigned (U.S. National Phase of PCT/GB99/00905)

Remarks

Claims 3-11, 13-16, 18-19, 21-22, 24-26, and 29 have been amended to delete multiple dependencies. The amendments do not add new matter and Applicants respectfully request that they be entered. Support for the amendments can be found throughout the specification (e.g. in the original claims). Upon entry of the forgoing amendment, claims 1-29 are pending in the application with claims 1, 17 and 27 being the independent claims.

It is not believed that extensions of time are required, beyond those that may otherwise be provided for in accompanying documents. However, if additional extensions of time are necessary to prevent abandonment of this application, then such extensions of time are hereby petitioned under 37 C.F.R. § 1.136(a), and any fees required therefor are hereby authorized to be charged to our Deposit Account No. 19-0036.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

Eric K. Steffe

Attorney for Applicants Registration No. 36,688

Date:

1100 New York Avenue, N.W.

Suite 600

Washington, D.C. 20005

(202) 371-2600

534 Rec'd PCT/PT3 20 SEP 2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Wyatt et al.

Appl. No.

For:

To be assigned (U.S.

National Phase of

PCT/GB99/00905)

Filed: Herewith

Signal Transduction Protein

Involved in Plant Dehiscence

Art Unit:

To be assigned

Examiner:

To be assigned

Atty. Docket: 0623.0890000/EKS/PSC

Claim For Priority Under 35 U.S.C. § 119(a)-(d) In Utility Application

Commissioner for Patents Washington, D.C. 20231

Sir:

Priority under 35 U.S.C. § 119(a)-(d) is hereby claimed to the following priority document(s), filed in a foreign country within twelve (12) months prior to the filing of the above-referenced United States utility patent application:

Country	Priority Document Appl. No.	Filing Date
Great Britain	GB9806113.8	20 March 1998

A certified copy of each listed priority document is submitted herewith. Prompt acknowledgment of this claim and submission is respectfully requested.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

Eric K. Steffe

Attorney for Applicants Registration No. 36,688

Suite 600

Washington, D.C. 20005-3934

(202) 371-2600

534 Rec'd PCT/PTO 20 SEP 2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Wyatt et al.

Appl. No. To be assigned

(U.S. Natl. Phase of PCT/GB99/00905)

Filed: Herewith

For: Signal Protein Involved in Plant

Dehiscence

Art Unit:

To be assigned

Examiner:

To be assigned

Atty. Docket: 0623.0890000/EKS/PSC

Preliminary Amendment and Submission of Sequence Listing

Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

In advance of prosecution, please amend the application as follows:

In the Specification:

Please insert the sequence listing at the end of the application.

On page 4, line 24, after "figure 1", please insert --(SEQ ID NO: 15)--.

On page 5, line 10, after "Figure 1", please insert -- (SEQ ID NO: 15)--.

On page 6, line 25, after "figure 1", please insert -- (SEQ ID NO: 15)--.

On page 8, line 17, after "figure 1", please insert --(SEQ ID NO: 14)--.

On page 8, line 19, after "Figure 1", please insert -- (SEQ ID NO: 14)--.

On page 9, line 22, after "figure 1", please insert -- (SEQ ID NO: 14)--.

On page 12, line 10, after "figure 1", please insert -- (SEQ ID NO: 15)--.

On page 12, line 22, after "Figure 1", please insert -- (SEQ ID NO: 15)--.

On page 14, line 12, after "Fig 5", please insert -- (SEQ ID NOs: 3, 5, and 6)--.

On page 14, line 12, after "Fig 9", please insert -- (SEQ ID NOs: 37 and 38)--.

On page 14, line 12, after "Fig 11", please insert -- (SEQ ID NOs: 7 and 8)--.

On page 18, line 15, after "Figure 15)", please insert -- (SEQ ID NOs: 31 and 32)--.

On page 18, line 16, after "16)", please insert -- (SEQ ID NOs: 33 and 34)--.

On page 18, line 16, after "(Figure 17)", please insert --(SEQ ID NOs: 35 and 36)--.

On page 19, line 9, after "Sac66", please insert --(SEQ ID NO: 31)--.

```
On page 19, line 9, after "DZ15", please insert --(SEQ ID NO: 33)--.
```

On page 19, line 10, after "OSR(7)", please insert --(SEQ ID NO: 35)--.

On page 19, line 22, after "sites", please insert --(SEQ ID NOs: 14 and 15)--.

On page 20, line 7, after "DZ2B", please insert --(SEQ ID NO: 23)--.

On page 20, line 16, after "DZ2AT3 peptide", please insert --(SEQ ID NOs: 27 and 28)--.

On page 20, line 20, after "DZ2AT3", please insert --(SEQ ID NO: 29)--.

On page 21, line 2, after "Sac66", please insert --(SEQ ID NOs: 31 and 32)--.

On page 21, line 4, after "DZ15", please insert --(SEQ ID NOs: 33 and 34)--.

On page 21, line 6, after "OSR7", please insert --(SEQ ID NOs: 35 and 36)--.

On page 22, line 22, after "TGG-3')", please insert --(SEQ ID NO: 1)--.

On page 23, line 1, after "TGG-3')", please insert --(SEQ ID NO: 1)--.

On page 23, line 1, after "CGA A -3"), please insert --(SEQ ID NO: 2)--.

On page 23, line 14, after "TGG-3")", please insert --(SEQ ID NO: 1)--.

On page 23, line 14, after "CGA A -3')", please insert --(SEQ ID NO: 2)--.

On page 23, line 18, after "(Figure 1)", please insert -- (SEQ ID NO: 14)--.

On page 24, line 3, after "(Figure 1)", please insert -- (SEQ ID NO: 14)--.

On page 25, line 15, after the first "(Figure 5)", please insert --(SEQ ID NO: 23)--.

On page 25, line 18, after "DZ2BFL", please insert -- (SEQ ID NO: 3)--.

On page 25, line 19, after "T7", please insert -- (SEQ ID NO: 4)--.

On page 26, line 13, after "DZ2BGENF", please insert -- (SEQ ID NO: 5)--.

On page 26, line 15, after "DZ2BGENR", please insert -- (SEQ ID NO: 6)--.

On page 27, line 7, after "Figure 9", please insert -- (SEQ ID NO: 27)--.

On page 27, line 28, after "(Figure 11)", please insert -- (SEQ ID NO: 29)--.

On page 28, line 1, after "ATDZ2F", please insert -- (SEQ ID NO: 7)--.

On page 28, line 2, after "ATDZ2R", please insert -- (SEQ ID NO: 8)--.

On page 30, line 5, after "DZ2FLA", please insert -- (SEQ ID NO: 9)--.

On page 30, line 6, after "DZ2RLA", please insert -- (SEQ ID NO: 10)--.

On page 31, line 15, after "Figure 15)", please insert -- (SEQ ID NOs: 31 and 32)--.

On page 31, line 16, after "16)", please insert -- (SEQ ID NOs: 33 and 34)--.

On page 31, line 16, after "Figure 17", please insert -- (SEQ ID NOs: 35 and 36)--.

On page 32, line 8, after "Figure 16)", please insert -- (SEQ ID NO: 33)--.

On page 32, line 15, after "T7", please insert -- (SEQ ID NO: 4)--.

On page 32, line 16, after "DZ15RL", please insert -- (SEQ ID NO: 11)--.

On page 32, line 25, after "F1", please insert -- (SEQ ID NO: 12)--.

On page 32, line 26, after "RI", please insert -- (SEQ ID NO: 13)--.

Remarks

The specification has been amended to direct the entry of this sequence listing after the claims of the above identified application and to provide sequence identifiers in all instances where the specification discusses the sequences. See 37 C.F.R. § 1.821(d).

In accordance with 37 C.F.R. § 1.821(g), this submission includes no new matter. In accordance with 37 C.F.R. § 1.821(f), the paper copy of the Sequence Listing and the computer readable copy of the Sequence Listing submitted herewith in the above application are the same.

It is respectfully believed that this application is now in condition for examination. Early notice to this effect is respectfully requested.

Respectfully submitted,

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.

Eric K! Steffe

Attorney for Applicants Registration No. 36,688

1100 New York Avenue, N.W.

Suite 600

Washington, D.C. 20005-3934

(202) 371-2600 P:\USERS\PChoi\0623\089\Seq List Submission Pleading.wpd SKGF Rev. 12/30/99 mac

SEQUENCE LISSA Rec'd PCT/PTG 20 SEP2000

```
<110> Wyatt, Paul
     Roberts, Jeremy A.
      Whitelaw, Catherine
 <120> Signal Transduction Protein Involved in Plant Dehiscence
<130> 0623.0890000
<140> To Be Assigned
<141> Herewith
<150> GB9806113.8
<151> 1998-03-20
<160> 38
<170> PatentIn Ver. 2.0
<210> 1
<211> 14
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: oligo dT primer 7
<400> 1
ttttttttt ttgg
                                                                    14
<210> 2
<211> 10
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: Arbitrary primer A
<400> 2
agccagcgaa
                                                                    10
<210> 3
<211> 19
<212> DNA
<213> Artificial Sequence
<220>
```

<223> Description of Artificial Sequence: Primer DZ2BFL	
<400> 3	
	19
aaccaagtca gtagaagtg	10
<210> 4	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: Primer T7	
<400> 4	18
aatacgactc actatagg	10
<210> 5	
<211> 26	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: Primer DZ2BGENF	
<400> 5 ggctctagac gaactgcgga gcaagg	26
ggetetagae gaaetgegga geaagg	
<210> 6	
<211> 29	
<212> DNA	
<213> Artificial Sequence	
.224	
<pre><220> <223> Description of Artificial Sequence: Primer DZ2BGENR</pre>	
22237 Description of Artificial bequence. Frimer bazbonik	
<400> 6	
ctgccatggt cggttttttt tcttcgaac	29
<210> 7	
<211> 23	
<212> DNA	
<213> Artificial Sequence	
<220>	
<pre><223> Description of Artificial Sequence: Primer ATDZ2F</pre>	
<u> </u>	
<400> 7	
cactagtagg gcacgcgtgg tcg	23
-210> 0	

<210> 8

```
<211> 27
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: Primer ATDZ2R
<400> 8
tccatggtcg atttcttttc tctcaag
                                                                    27
<210> 9
<211> 28
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Primer DZ2FLA
<400> 9
ggcgaattcc ggtgaggagg cagtaatc
                                                                    28
<210> 10
<211> 28
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Primer DZ2RLA
<400> 10
ggcccatggc atacatacac acttagac
                                                                    28
<210> 11
<211> 22
<212> DNA
<213> Artificial Sequence
<223> Description of Artificial Sequence: Primer DZ15RL
<400> 11
aacagctgaa aacctcacga ag
                                                                    22
<210> 12
<211> 28
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Primer F1
```

<400> 12 ggcccatggc tgccaagctt tgagtagc	28
<210> 13 <211> 27 <212> DNA	
<213> Artificial Sequence	
<220> <223> Description of Artificial Sequence: Primer R1	
<400> 13 ggcctgcagt gcctaggatc tggaagc	27
<210> 14 <211> 605 <212> DNA <213> Brassica napus	
<220> <221> CDS <222> (20)(427)	
<pre><400> 14 ggcacgagca gaatcgaag atg gca aca aaa tcc atg gga gat atc gag aaa</pre>	52
ata aag aag aaa cta aac gtg ttg atc gtc gat gat gat cca cta aac Ile Lys Lys Lys Leu Asn Val Leu Ile Val Asp Asp Asp Pro Leu Asn 15 20 25	100
ctt ata att cat gag aag atc atc aaa gcg att ggg ggt att tca cag Leu Ile Ile His Glu Lys Ile Ile Lys Ala Ile Gly Gly Ile Ser Gln 30 35 40	148
aca gcg aat aac ggt gag gag gca gta atc atc cac cgt gac ggc ggc Thr Ala Asn Asn Gly Glu Glu Ala Val Ile Ile His Arg Asp Gly Gly	196
45 50 55	
tca tct ttt gac ctt atc cta atg gat aaa gaa atg ccc gag agg gat Ser Ser Phe Asp Leu Ile Leu Met Asp Lys Glu Met Pro Glu Arg Asp 60 65 70 75	244
tca tct ttt gac ctt atc cta atg gat aaa gaa atg ccc gag agg gat Ser Ser Phe Asp Leu Ile Leu Met Asp Lys Glu Met Pro Glu Arg Asp	244

<400> 16

ttc atg Phe Met															388
gac aag Asp Lys 125												tgai	tggai	tat	437
atatttt	ata t	tate	gaaa	ac ac	cacat	taata	a ac	gtct	aagt	gtg	tatg	tat 🤉	gcata	agatac	497
ttgcatg	tgt g	tgtt	ttaç	ja at	ttaç	gggtt	c ct	ttat	cgtc	cgt	gatat	tat a	aatca	atgtaa	557
gttgttg	ctt t	aago	ttat	ta aa	aatat	cttaa	a ata	aagg	gttt	cct	ctaco	C			605
<210> 1! <211> 1: <212> PP <213> Bi	36 RT	ca n	ıapus	3											
<400> 19 Met Ala 1		Lys	Ser 5	Met	Gly	Asp	Ile	Glu 10	Lys	Ile	Lys	Lys	Lys 15	Leu	
Asn Val	Leu	Ile 20	Val	Asp	Asp	Asp	Pro 25	Leu	Asn	Leu	Ile	Ile 30	His	Glu	
Lys Ile	Ile 35	Lys	Ala	Ile	Gly	Gly 40	Ile	Ser	Gln	Thr	Ala 45	Asn	Asn	Gly	
Glu Glu 50	Ala	Val	Ile	Ile	His 55	Arg	Asp	Gly	Gly	Ser 60	Ser	Phe	Asp	Leu	
Ile Leu 65	Met :	Asp	Lys	Glu 70	Met	Pro	Glu	Arg	Asp 75	Gly	Val	Ser	Thr	Thr 80	
Lys Lys	Leu 2	Arg	Glu 85	Met	Glu	Val	Lys	Ser 90	Met	Ile	Val	Gly	Val 95	Thr	
Ser Leu		Asp 100	Asn	Glu	Glu	Glu	Arg 105	Arg	Ala	Phe	Met	Glu 110	Ala	Gly	
Leu Asn	His (Cys	Leu	Ala	Lys	Pro 120	Leu	Thr	Lys	Asp	Lys 125	Ile	Ile	Pro	
Leu Ile 130	Asn (Gln	Leu	Met	Asp 135	Ala									
<210> 16	5														
<211> 13															
<212> PR		ca n	apus	}											
<7T\$> BI	assi	ca n	apus	;											

Met Ala Thr Lys Ser Met Gly Asp Ile Glu Lys Ile Lys Lys Leu 1 5 10 15

Asn Val Leu Ile Val Asp Asp Asp Pro Leu Asn Leu Ile Ile His Glu 20 25 30

Lys Ile Ile Lys Ala Ile Gly Gly Ile Ser Gln Thr Ala Asn Asn Gly
35 40 45

Glu Glu Ala Val Ile Ile His Arg Asp Gly Gly Ser Ser Phe Asp Leu
50 55 60

Ile Leu Met Asp Lys Glu Met Pro Glu Arg Asp Gly Val Ser Thr Thr 65 70 75 80

Lys Lys Leu Arg Glu Met Glu Val Lys Ser Met Ile Val Gly Val Thr
85 90 95

Ser Leu Ala Asp Asn Glu Glu Glu Arg Arg Ala Phe Met Glu Ala Gly 100 105 110

Leu Asn His Cys Leu Ala Lys Pro Leu Thr Lys Asp Lys Ile Ile Pro 115 120 125

Leu Ile Asn Gln Leu Met Asp Ala 130 135

<210> 17

<211> 132

<212> PRT

<213> Escherichia coli

<400> 17

Met Gln Glu Asn Tyr Lys Ile Leu Val Val Asp Asp Asp Met Arg Leu 1 5 10 15

Arg Ala Leu Leu Glu Arg Tyr Leu Thr Glu Gln Gly Phe Gln Val Arg 20 25 30

Ser Val Ala Asn Ala Glu Gln Met Asp Arg Leu Leu Thr Arg Glu Ser 35 40 45

Phe His Leu Met Val Leu Asp Leu Met Leu Pro Gly Glu Asp Gly Leu 50 55 60

Ser Ile Cys Arg Arg Leu Arg Ser Gln Ser Asn Pro Met Pro Ile Ile 65 70 75 80

Met Val Thr Ala Lys Gly Glu Glu Val Asp Arg Ile Val Gly Leu Glu 85 90 95

Ile Gly Ala Asp Asp Tyr Ile Pro Lys Pro Phe Asn Pro Arg Glu Leu 100 105 110

Leu Ala Arg Ile Arg Ala Val Leu Arg Arg Gln Ala Asn Glu Leu Pro 115 120 125

Gly Ala Pro Ser 130

<210> 18

<211> 126

<212> PRT

<213> Escherichia coli

<400> 18

Met Ala Arg Arg Ile Leu Val Val Glu Asp Glu Ala Pro Ile Arg Glu

1 10 15

Met Val Cys Phe Val Leu Glu Gln Asn Gly Phe Gln Pro Val Glu Ala 20 25 30

Glu Asp Tyr Asp Ser Ala Val Asn Gln Leu Asn Glu Pro Trp Pro Asp 35 40 45

Leu Ile Leu Leu Asp Trp Met Leu Pro Gly Gly Ser Gly Ile Gln Phe 50 55 60

Ile Lys His Leu Lys Arg Glu Ser Met Thr Arg Asp Ile Pro Val Val 65 70 75 80

Met Leu Thr Ala Arg Gly Glu Glu Glu Asp Arg Val Arg Gly Leu Glu 85 90 95

Thr Gly Ala Asp Asp Tyr Ile Thr Lys Pro Phe Ser Pro Lys Glu Leu 100 105 110

Val Ala Arg Ile Lys Ala Val Met Arg Arg Ile Ser Pro Met 115 120 125

<210> 19

<211> 144

<212> PRT

<213> Salmonella typhimurium

<400> 19

Met Gln Arg Gly Ile Val Trp Val Val Asp Asp Asp Ser Ser Ile Arg 1 5 10 15

Trp Val Leu Glu Arg Ala Leu Ala Gly Ala Gly Leu Thr Cys Thr Thr 20 25 30

Phe Glu Asn Gly Asn Asn Thr Arg Cys Glu Val Leu Ala Ala Leu Ala 35 40 45

Ser Lys Thr Pro Asp Val Leu Leu Ser Asp Ile Arg Met Pro Gly Met 50 55 60

Asp Gly Leu Ala Leu Leu Lys Gln Ile Lys Gln Arg His Pro Met Leu 65 70 75 80

Pro Val Ile Ile Met Thr Ala Asn Thr Arg Cys His Ser Asp Leu Asp 85 90 95

Ala Ala Val Ser Ala Tyr Gln Gln Gly Ala Phe Asp Tyr Leu Pro Lys
100 105 110

Pro Phe Asp Ile Asp Glu Ala Val Ala Leu Val Glu Arg Ala Ile Ser 115 120 125

His Tyr Gln Glu Gln Gln Pro Arg Asn Ile Glu Val Asn Gly Pro 130 135 140

<210> 20

<211> 124

<212> PRT

<213> Bacillus subtilis

<400> 20

Met Met Asn Glu Lys Ile Leu Ile Val Asp Asp Gln Tyr Gly Ile Arg 1 5 10 15

Ile Leu Leu Asn Glu Val Phe Asn Lys Glu Gly Tyr Gln Thr Phe Gln 20 25 30

Ala Ala Asn Gly Leu Gln Ala Leu Asp Ile Val Thr Lys Glu Arg Pro 35 40 45

Asp Leu Val Leu Leu Asp Met Lys Ile Pro Gly Met Asp Gly Ile Glu 50 60

Ile Leu Lys Arg Met Lys Val Ile Asp Glu Asn Ile Arg Val Ile Ile 65 70 75 80

Met Thr Ala Tyr Gly Glu Leu Asp Met Ile Gln Glu Ser Lys Glu Leu 85 90 95

Gly Ala Leu Thr His Phe Ala Lys Pro Phe Asp Ile Asp Glu Ile Arg

Asp Ala Val Lys Lys Tyr Leu Pro Leu Lys Ser Asn 115 120

<210> 21

<211> 129

<212> PRT

<213> Escherichia coli

<400> 21

Met Ala Asp Lys Glu Leu Lys Phe Leu Val Val Asp Asp Phe Ser Thr

Met Arg Arg	Ile Val	Arg Asn	Leu Leu 25	Lys Glu	Leu	Gly	Phe 30	Asn	Asn
Val Glu Glu 35		Asp Gly	Val Asp 40	Ala Leu	Asn	Lys 45	Leu	Gln	Ala
Gly Gly Tyr 50	Gly Phe	Val Ile 55	Ser Asp	Trp Asn	Met 60	Pro	Asn	Met	Asp
Gly Leu Glu 65	Leu Leu	Lys Thr 70	Ile Arg	Ala Asp		Ala	Met	Ser	Ala 80
Leu Pro Val	Leu Met 85	Val Thr	Ala Glu	Ala Lys 90	Lys	Glu	Asn	Ile 95	Ile
Ala Ala Ala	Gln Ala	Gly Ala	Ser Gly 105	Tyr Val	Val	Lys	Pro 110	Phe	Thr
Pro Ala Thr		Glu Lys	Leu Asn 120	Lys Ile	Phe	Glu 125	Lys	Leu	Gly
Met									
<210> 22 <211> 111 <212> PRT <213> Arabi	dopsis t	haliana							
<220> <221> Unsur <222> 67 <223> Xaa=		o acid							
<400> 22 Leu Lys Val	Leu Val		Glu Asn	Gly Val	. Ser	Arg	Met	Val 15	Thr
Lys Gly Leu	Leu Val	His Lev	Gly Cys 25	Glu Val	Thr	Thr	Val 30	Ser	Ser
Asn Glu Glu 35	_	Arg Val	Val Ser 40	His Glu	His	Lys 45	Val	Val	Phe
Met Asp Val 50	. Cys Met	Pro Gly		Asn Tyr	Gln 60	Ile	Ala	Leu	Arg
Ile His Xaa	Pro Leu	_	Ala Leu	Ser Gly		Thr	Asp	Lys	Ser 80
		70		, 3					

Pro Val Ser Leu Asp Asn Ile Arg Asp Val Leu Ser Asp Leu Leu
100 105 110

```
<210> 23
<211> 1716
<212> DNA
<213> Brassica napus
<220>
<221> CDS
<222> (1516)..(1716)
<220>
<221> Unsure
<222> 48
<223> n= any nucleotide
<220>
<221> Unsure
<222> 56
<223> n= any nucleotide
<220>
<221> Unsure
<222> 57
<223> n= any nucleotide
<220>
<221> Unsure
<222> 62
<223> n= any nucleotide
<220>
<221> Unsure
<222> 72
<223> n= any nucleotide
<220>
<221> Unsure
<222> 76
<223> n= any nucleotide
<220>
<221> Unsure
<222> 80
<223> n= any nucleotide
<220>
<221> Unsure
<222> 103
<223> n= any nucleotide
<220>
<221> Unsure
```

```
<222> 137
<223> n= any nucleotide
<220>
<221> Unsure
<222> 146
<223> n= any nucleotide
<220>
<221> Unsure
<222> 599
<223> n= any nucleotide
<220>
<221> Unsure
<222> 602
<223> n= any nucleotide
<220>
<221> Unsure
<222> 614
<223> n= any nucleotide
<220>
<221> Unsure
<222> 626
<223> n= any nucleotide
<220>
<221> Unsure
<222> 627
<223> n= any nucleotide
<220>
<221> Unsure
<222> 673
<223> n= any nucleotide
tatataaata cggtttaaca gatatgttct ggttataaat gtaattcnat gtgccnntca 60
anttttattt tnattngttn tactagggac attagtttta acnttttata tatcatgtaa 120
caaaaaaaaa aaaaacnttt tatatntcaa ctatgagcaa ttattcttat agtgttttct 180
ttttccagaa atttgacgac aacctaacta aaacaattta atttgacgtt agttaagtaa 240
tttatataga tggataaatt gagcaagcac attacgaact gcggatcaag gagagtcaca 300
atttaattct tacgttatac acaaaattat ctaaatacta tatatatata cagctgcatg 360
ctacgataat gatcaaatgt ttatgtactt ttcagcgaaa attcttgtcg ccatacatta 420
ctgtgttaat gaatcattaa atatgtgaag gaggaaaaga gtacaaaagg agttttgttg 480
```

aggcatttcg	cagacactga	aatgtgaata	ataataaagg	aattgccgaa	ttgatttcta	540
gttggtgaag	tgggtgaaaa	ttgtatgtcc	attgcttata	aactataaaa	tataatatnt	600
tnatattatc	actntggaca	ttagtnngat	agaccctagc	taaaattttt	aaaaattata	660
cattcatttt	ctnaagtacc	aaacttaatt	atcacaatcg	gataaaattg	tttaagaaac	720
cattacaaac	tcagcttgtg	gactctgaga	gaaactaaga	gctagacata	cggttagtag	780
tgtagccgca	ttttttatgc	ttaatttgct	taagcatgac	ttctatgctc	cttgatgata	840
tttattttaa	tatcctagga	catatggatt	tgataaagat	cttatcaacc	tttcaacaag	900
accattagct	caacaaacaa	aatactgaaa	gtatataatc	ttggttacag	aattcttatg	960
ccaaaaatat	cataatatat	atagaattcg	gttatgatta	agatgaatta	tttaattaat	1020
atatttttca	cttttgtttt	cttatgtatt	cttagtattt	gttcaccata	ttgaccgatt	1080
ggtgtcatat	tagtttggta	agacaactca	gttgcaacga	tgcagattac	atttcaggaa	1140
gattcatgta	agaaagatat	ttcgctttgt	ggtgtgaaaa	tatgcctctt	tcactttttt	1200
tcaactataa	atttcgatcg	atgtatctac	gttcttaaca	caattcacaa	tcttctttag	1260
aatccaaaat	tgtaagccgc	tttctaatct	ctttctcagt	atacatatgt	aatatgtatg	1320
catatattat	tattcataat	acaaacacga	acccatgcat	gcaagaagat	agttacacgc	1380
tcataacaaa	cacaaaaaaa	catacgcatg	cattagaaca	cttgtatgtt	aatttccata	1440
atgttttgca	taaacattct	tcgttttaat	tagcttcttt	ttgtgtgaag	attgttcgaa	1500
gaaaaaaaac	cgaag atg Met 1	gca aca acg Ala Thr Thr	tca aca tc Ser Thr Se 5	r Thr Gly A	at atc aag sp Ile Lys 10	1551
aaa acc aa Lys Thr Ly 1	s Ser Val G	aa gtg aag lu Val Lys 20	aag aaa ctt Lys Lys Leu	aac gtg tt Asn Val Le 25	g atc gtt u Ile Val	1599
gat gat ga Asp Asp As 30	t aca gta a p Thr Val I	tt cgt aaa le Arg Lys 35	ctt cac gag Leu His Glu	aat atc at Asn Ile Il 40	c aaa tcg e Lys Ser	1647
atc ggt gg Ile Gly Gl 45	a att tca c y Ile Ser G	ag acg gct In Thr Ala	aag aac ggt Lys Asn Gly 55	gag gag gc Glu Glu Al	a gtg aac a Val Asn 60	1695
_	c gac ggc a g Asp Gly <i>I</i> 65					1716

```
<210> 24
<211> 67
<212> PRT
<213> Brassica napus
<400> 24
Met Ala Thr Thr Ser Thr Ser Thr Gly Asp Ile Lys Lys Thr Lys Ser
Val Glu Val Lys Lys Leu Asn Val Leu Ile Val Asp Asp Asp Thr
                                  25
Val Ile Arg Lys Leu His Glu Asn Ile Ile Lys Ser Ile Gly Gly Ile
Ser Gln Thr Ala Lys Asn Gly Glu Glu Ala Val Asn Ile His Arq Asp
     50
Gly Asn Ala
 65
<210> 25
<211> 576
<212> DNA
<213> Brassica napus
<220>
<221> Unsure
<222> 6
<223> n= any nucleotide
<220>
<221> Unsure
<222> 518
<223> n= any nucleotide
<400> 25
tegtenatga tgateetgta ataegtaaac tteaegagat tateateaaa teaateqqtq 60
gaatttcaca gacagctaag aacggtgagg aggcagtgaa catccaccgc gacggcaatg 120
catctttcga ccttatccta atggataaag aaatgcccga gagggatgga ctttcggcaa 180
ctaagaagct aagagaaatg aaagtgacgt ctatgattat tggggtgacg acactggctg 240
acaatgaaga ggaacgtaag gctttcatgg aagctggact taaccattgc ttggcaaaac 300
ccttaagcaa agccaagatc ctccctctca tcaacaatct catggatgct tgatggatgg 360
atgaattgtc gccactacat atctacatta tacaaatatg aaaaacacat ataataacgt 420
catacacctg tgtgtgtatg catagatatc tatccgcatg tgtgttttta gggttgttat 480
```

gtttgatttt tattgtgcgt ggcgtgatat acaatcangt nagtcgttac ttttggctta 540

taaaataatg aataagattt gttaaaaata aaaaaa

576

<210> 26

<211> 116

<212> PRT

<213> Brassica napus

<220>

<221> Unsure

<222> 2

<223> Xaa= any amino acid

<400> 26

Val Xaa Asp Asp Pro Val Ile Arg Lys Leu His Glu Ile Ile Ile Lys

1 10 15

Ser Ile Gly Gly Ile Ser Gln Thr Ala Lys Asn Gly Glu Glu Ala Val $20 \hspace{1.5cm} 25 \hspace{1.5cm} 30$

Asn Ile His Arg Asp Gly Asn Ala Ser Phe Asp Leu Ile Leu Met Asp 35 40 45

Lys Glu Met Pro Glu Arg Asp Gly Leu Ser Ala Thr Lys Lys Leu Arg 50 55 60

Glu Met Lys Val Thr Ser Met Ile Ile Gly Val Thr Thr Leu Ala Asp 65 70 75 80

Asn Glu Glu Glu Arg Lys Ala Phe Met Glu Ala Gly Leu Asn His Cys
85 90 95

Leu Ala Lys Pro Leu Ser Lys Ala Lys Ile Leu Pro Leu Ile Asn Asn 100 105 110

Leu Met Asp Ala 115

<210> 27

<211> 818

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (180)..(605)

<220>

<221> Unsure

<222> 350

<223> n= any nucleotide

<220>

<221> Unsure <222> 57 <223> Xaa= any amino acid <400> 27 atatatgtga tacagataca tctatataca aattaaacac gaaaccatac atgcacggtg 60 tgatcacaca cgcacacaca tagaaacata aacacgcaat aatttctata caqtttaatt 120 tcatttttaa cttacttctt tttttttggt gaagattctt gagagaaaag aaatcgaag atg gca aca aaa tcc acc gga ggt acc gag aaa acc aag tcg ata gaa 227 Met Ala Thr Lys Ser Thr Gly Gly Thr Glu Lys Thr Lys Ser Ile Glu gtg aag aag aaa cta atc aac gtg ttg atc gtc gat gat gat cca tta 275 Val Lys Lys Leu Ile Asn Val Leu Ile Val Asp Asp Pro Leu aac cgt aga ctc cac gag atg atc atc aaa acg atc gga gga att tct 323 Asn Arg Arg Leu His Glu Met Ile Ile Lys Thr Ile Gly Gly Ile Ser cag act gca aag aat ggc gaa gag gcn gtg atc ctc cac cgt gac ggc Gln Thr Ala Lys Asn Gly Glu Glu Xaa Val Ile Leu His Arg Asp Gly 50 55 gaa gca tot tto gac ott att ota atg gat aag gaa atg oot gag agg 419 Glu Ala Ser Phe Asp Leu Ile Leu Met Asp Lys Glu Met Pro Glu Arg 65 70 gat gga gtt tcg aca att aag ang cta aga gaa atg aaa ggg acg tca 467 Asp Gly Val Ser Thr Ile Lys Xaa Leu Arg Glu Met Lys Gly Thr Ser 85 95 atg atc gtt ggg gta acg tca gta gct gac caa gaa gaa gag cgt aag 515 Met Ile Val Gly Val Thr Ser Val Ala Asp Gln Glu Glu Glu Arg Lys 100 110 get tit atg gaa get ggg ete aac eat tge ttg gaa aaa eee tta ace 563 Ala Phe Met Glu Ala Gly Leu Asn His Cys Leu Glu Lys Pro Leu Thr 120 aag gcc aag atc ttc ccg ctc att agc cac ctc ttc gat gct 605 Lys Ala Lys Ile Phe Pro Leu Ile Ser His Leu Phe Asp Ala 135 tgatggatga aggeteatta atgtatetat atttteaate atgaaateae etacaegtgt 665 atttgacaca aaaatctgca tttgttgtga tatagggttt ctcatatcta tgtttgattt 725 attiticttat cgtccgaggt aaaatcatgc aagtcattic tittiggctaa taaaatatta 785 aaataaggtt ttctcaaaaa aaaaaaaaa aaa 818

<220>

<221> Unsure

```
<210> 28
<211> 142
<212> PRT
<213> Arabidopsis thaliana
<220>
<221> Unsure
<222> 57
<223> Xaa= any amino acid
<220>
<221> Unsure
<222> 88
<223> Xaa= any amino acid
<400> 28
Met Ala Thr Lys Ser Thr Gly Gly Thr Glu Lys Thr Lys Ser Ile Glu
Val Lys Lys Leu Ile Asn Val Leu Ile Val Asp Asp Pro Leu
             20
                                                      30
Asn Arg Arg Leu His Glu Met Ile Ile Lys Thr Ile Gly Gly Ile Ser
Gln Thr Ala Lys Asn Gly Glu Glu Xaa Val Ile Leu His Arg Asp Gly
                         55
Glu Ala Ser Phe Asp Leu Ile Leu Met Asp Lys Glu Met Pro Glu Arg
Asp Gly Val Ser Thr Ile Lys Xaa Leu Arg Glu Met Lys Gly Thr Ser
Met Ile Val Gly Val Thr Ser Val Ala Asp Gln Glu Glu Glu Arg Lys
            100
                                105
Ala Phe Met Glu Ala Gly Leu Asn His Cys Leu Glu Lys Pro Leu Thr
                            120
        115
Lys Ala Lys Ile Phe Pro Leu Ile Ser His Leu Phe Asp Ala
                        135
<210> 29
<211> 1324
<212> DNA
<213> Arabidopsis thaliana
<220>
<221> CDS
<222> (1200)..(1322)
```

<222> 1142 <223> n= any nucleotide

<400> 29 qtaatqcqac tcactataqq qcacqcqtgg tcgacqgccc gggctggtcc tcattcgtat 60 ttaacatqta tqtqqqatat ttqqctataa attatgtaaa aaatttcacg atagattgtt 180 gaattttttga atttcgagtt aaaatatctt caaattacct cacatttaca aaaaggtaga 240 actgttgaaa aactaatget ctataaaaca ctagacaata acaaaatacg taatgegtaa 300 agaacctaaa ttatgatttt atttatcttt cttccttttt ccgtgagtat aagccatttt 360 tcatagtaaa gcattacgaa tacgacattg aacactactg acatataaag tagtagattt 420 tgatgggtta acttgtatgc ttaatttgct taagcatgaa cttcaatgct tttataaaag 480 tacttcatqa qaatattcct cqttctatac tagcagaagg gttcgatagt gattttacaa 540 ccgttcaaca aaacctttaa acccaaaaaa ccaaagaatg aaagtatcta aacttgatta 600 tacatttctt gtctaaatta tcaaataaca tactctcttt tgtttactta taaacgatat 660 gaaagaaata aataaaaaga acatagaatc ttattatgat ctagaagaat taattaaaga 720 aatatatata talttttttt catttctact catgtttctt atacattctt taaatttgtt 780 caccattttg atttacttgt tctcatatta gtttgttata caactcactt agaataatgt 840 agattacatt tcagccaaat tcatgtaaaa gatgcttttc tttgtgatgt ttttaaaatg 900 ctttcttttc acttttttc tttcttaact ataaatcttg atcgaatgcc taccttctta 960 qaacataaqa tottotttaa aatocaaaat ogtaggocac tatttoatta taottatgta 1020 atatatgtga tacagataca tntatataca aattaaacac gaaaccatac atgcacggtg 1080 tgatcacaca cgcacacaca tagaaacata aacacgcaat aatttctata cagtttaatt 1140 tcatttttaa cttacttett tttttttggt gaagattett gagagaaaag aaategaag 1199 atg gca aca aaa tcc acc gga ggt acc gag aaa acc aag tcg ata gaa 1247 Met Ala Thr Lys Ser Thr Gly Gly Thr Glu Lys Thr Lys Ser Ile Glu 1 5 15 gtg aag aag aaa cta atc aac gtg ttg atc gtc gat gat gat cca tta 1295 Val Lys Lys Lys Leu Ile Asn Val Leu Ile Val Asp Asp Asp Pro Leu 20 1324 aac cgt aga ctc cac gag tgt cat caa aa Asn Arg Arg Leu His Glu Cys His Gln

35

<210> 30

<211> 41 <212> PRT <213> Arabidopsis thaliana <400> 30 Met Ala Thr Lys Ser Thr Gly Gly Thr Glu Lys Thr Lys Ser Ile Glu Val Lys Lys Leu Ile Asn Val Leu Ile Val Asp Asp Pro Leu 25 Asn Arg Arg Leu His Glu Cys His Gln <210> 31 <211> 1657 <212> DNA <213> Brassica napus <220> <221> CDS <222> (145)..(1443) <400> 31 ggcatcacga gggtacccgt aaatcccacc atacaacaaa gttctgtgaa agtctcccaa 60 aaactgcaaa gagtctcata ttagttctta ctctcagaaa taaaacacac tctttctgaa 120 aagattageg tttcaaaccc egaa atg gee egt tgt cat gga agt ett get 171 Met Ala Arg Cys His Gly Ser Leu Ala att ttc tta tqc qtt ctt ttg atg ctc gct tgc tgc caa gct ttg agt 219 Ile Phe Leu Cys Val Leu Leu Met Leu Ala Cys Cys Gln Ala Leu Ser 10 agc aac gta gat gga tat ggt cat gaa gat gga agc ttc gaa acc 267 Ser Asn Val Asp Asp Gly Tyr Gly His Glu Asp Gly Ser Phe Glu Thr gat agt tta atc aag ctc aac aac gac gac gac gtt ctt acc ttg aaa 315 Asp Ser Leu Ile Lys Leu Asn Asp Asp Asp Val Leu Thr Leu Lys 363 age tee gat aga eee act ace gaa tea tea act gtt agt gtt teg aac Ser Ser Asp Arg Pro Thr Thr Glu Ser Ser Thr Val Ser Val Ser Asn 60 ttc qqa qca aaa ggt gat gga aaa acc gat gat act cag gct ttc aag 411 Phe Gly Ala Lys Gly Asp Gly Lys Thr Asp Asp Thr Gln Ala Phe Lys 80 75 aaa gca tgg aag aag gca tgt tca aca aat gga gtg act act ttc ttg 459

Lys 90	Ala	Trp	Lys	Lys	Ala 95	Cys	Ser	Thr	Asn	Gly 100	Val	Thr	Thr	Phe	Leu 105	
	cct Pro															507
	tgc Cys															555
	aca Thr															603
	gag Glu 155															651
	gat Asp															699
	tct Ser															747
	aac Asn															795
	cag Gln															843
	atc Ile 235															891
	gct Ala															939
	gat Asp															987
	tta Leu															1035
	gac Asp															1083
acg	ctc	tct	gag	act	gac	aat	gga	gta	aga	atc	aag	act	tac	cag	gga	1131

Thr																
	Leu 315	Ser	Glu	Thr	Asp	Asn 320	Gly	Val	Arg	Ile	Lys 325	Thr	Tyr	Gln	Gly	
			act Thr													1179
			aat Asn													1227
			caa Gln 365													1275
			caa Gln													1323
			aaa Lys													1371
			gga Gly													1419
	200	att	tct	cct	aaa	tgc	cct	taai	ttac	taa g	gttga	attai	tg ta	aata	tacat	1473
		_	Ser		Lys	Cys	Pro									
Gly	Thr	Val	Ser	Pro 430	-	-		c ato	ctata	atcc	ttat	cta	egt :	attg	attctc	1533
Gly	Thr tacg	Val	Ser tata	Pro 430 tgtgg	- gt ta	ataga	atgc								attctc attgtt	
Gly aaa gata	Thr tacg	Val tat t	Ser tata gaaa	Pro 430 tgtgg	- gt ta ag ga	ataga attta	atgco	g aat	cata	cata	caat	agti	tga 🤉	gata		1593
Gly aaa gata	Thr tacg atata ttgt	Val tat t	Ser tata gaaa	Pro 430 tgtgg	- gt ta ag ga	ataga attta	atgco	g aat	cata	cata	caat	agti	tga 🤉	gata	attgtt	1593
Gly aaa gat gtc tgt <21 <21 <21	Thr tacgt atata ttgta C 0> 3: 1> 4: 2> Pi	Val	Ser tata gaaa	Pro 430 tgtgg actaa tcact	gt ta	ataga attta	atgco	g aat	cata	cata	caat	agti	tga 🤉	gata	attgtt	1593 1653
Gly aaa gat gtc tgt <21 <21 <21 <40	Thr tacgf atata ttgta c 0 > 3: 1 > 4: 2 > Pi 3 > B: 0 > 3:	Val tat f ata g ata f 2 33 RT rass:	Ser tata gaaaa tggt	Pro 430 tgtgg actas tcact	gt ta ag ga ag aa	ataga attta	atge atggg	g aat	tgtc	cata	caat	tagtt	gaa (gata	attgtt	1593 1653
Gly aaa gat gtc tgt <21 <21 <21 <40	Thr tacgf atata ttgta c 0 > 3: 1 > 4: 2 > Pi 3 > B: 0 > 3:	Val tat f ata g ata f 2 33 RT rass:	Ser tata gaaaa tggt	Pro 430 tgtgg actas tcact	gt ta ag ga ag aa	ataga attta	atge atggg	g aat	tgtc	cata	caat	tagtt	gaa (gata	attgtt gtcatt	1593 1653
gly aaa gat gtc tgt <21 <21 <21 <40 Met	Thr tacg atata ttgta 0 > 3: 1 > 4: 2 > P! 3 > B: 0 > 3: Ala	Val tat data data data data data data data	Ser tata gaaaa tggt	Pro 430 tgtgg actaa tcact	gt ta ag ga tg aa	ataga attta agtto Ser	atgco atggg gatto Leu	g aat	tgtco	cata cacg	caat	tagt†	gaa g	gata taat Leu 15	attgtt gtcatt Leu	1593 1653

Asn Asp Asp Val Leu Thr Leu Lys Ser Ser Asp Arg Pro Thr Thr

Glu Ser Ser Thr Val Ser Val Ser Asn Phe Gly Ala Lys Gly Asp Gly

Ile Lys Phe Gln Asn Ile Arg Met Asp Asn Val Lys Asn Pro Ile Ile 345

325

Ile Asp Gln Asn Tyr Cys Asp Lys Asp Lys Cys Glu Gln Gln Glu Ser 355 360 365	
Ala Val Gln Val Asn Asn Val Val Tyr Arg Asn Ile Gln Gly Thr Ser 370 375 380	
Ala Thr Asp Val Ala Ile Met Phe Asn Cys Ser Val Lys Tyr Pro Cys 385 390 395 400	
Gln Gly Ile Val Leu Glu Asn Val Asn Ile Lys Gly Gly Lys Ala Ser 405 410 415	
Cys Lys Asn Val Asn Val Lys Asp Lys Gly Thr Val Ser Pro Lys Cys 420 425 430	
Pro	
<210> 33 <211> 569 <212> DNA <213> Brassica napus	
<220> <221> CDS <222> (3)(311)	
<400> 33	
ag gtg acc gtt gct gat ggc aat gtg ctg gtc aag cga gag gta gac Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp 1 5 10	47
Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp	47 95
Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp 1 5 10 15 ggt ggc ttg gag aca gtt aaa gtc aaa ttg cca gct gtc att agc gcc Gly Gly Leu Glu Thr Val Lys Val Lys Leu Pro Ala Val Ile Ser Ala	
Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp 1 5 10 15 ggt ggc ttg gag aca gtt aaa gtc aaa ttg cca gct gtc att agc gcc Gly Gly Leu Glu Thr Val Lys Val Lys Leu Pro Ala Val Ile Ser Ala 20 25 30 gac ttg cgg ctc aat gag ccg cgg tac gct act ctg ccc aat atc atg Asp Leu Arg Leu Asn Glu Pro Arg Tyr Ala Thr Leu Pro Asn Ile Met	95
Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp 1 5 10 15 ggt ggc ttg gag aca gtt aaa gtc aaa ttg cca gct gtc att agc gcc Gly Gly Leu Glu Thr Val Lys Val Lys Leu Pro Ala Val Ile Ser Ala 20 25 30 gac ttg cgg ctc aat gag ccg cgg tac gct act ctg ccc aat atc atg Asp Leu Arg Leu Asn Glu Pro Arg Tyr Ala Thr Leu Pro Asn Ile Met 35 40 45 aag gcc aag aag aag ccc atc aaa aag ctc aca gcc aca gat gtc ggt Lys Ala Lys Lys Lys Pro Ile Lys Lys Leu Thr Ala Thr Asp Val Gly	95 143
Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp 1 5 10 15 ggt ggc ttg gag aca gtt aaa gtc aaa ttg cca gct gtc att agc gcc Gly Gly Leu Glu Thr Val Lys Val Lys Leu Pro Ala Val Ile Ser Ala 20 25 30 gac ttg cgg ctc aat gag ccg cgg tac gct act ctg ccc aat atc atg Asp Leu Arg Leu Asn Glu Pro Arg Tyr Ala Thr Leu Pro Asn Ile Met 35 40 45 aag gcc aag aag aag ccc atc aaa aag ctc aca gcc aca gat gtc ggt Lys Ala Lys Lys Lys Pro Ile Lys Lys Leu Thr Ala Thr Asp Val Gly 50 55 60 gtg gac ttg gcg cca cgt caa caa gtg ttg agc gta gaa gac ccg ccc Val Asp Leu Ala Pro Arg Gln Gln Val Leu Ser Val Glu Asp Pro Pro	95 143 191

gttcttacaa attcttcgtg aggttttcag ctgttaccaa taatatttt tcaaaatcga 401
ttttatttta cttgtaattt aaaagatcaa atattaatac aatgaacatt tttgtaacag 461
caatcttttt tttatatttt ggagatttca tcgacttatg tcataattat ttttatcaat 521
ttattgttgt ttgttagtga tataataaag tatatttct ggtcaaaa 569

<210> 34

<211> 103

<212> PRT

<213> Brassica napus

<400> 34

Val Thr Val Ala Asp Gly Asn Val Leu Val Lys Arg Glu Val Asp Gly

1 10 15

Gly Leu Glu Thr Val Lys Val Lys Leu Pro Ala Val Ile Ser Ala Asp 20 25 30

Leu Arg Leu Asn Glu Pro Arg Tyr Ala Thr Leu Pro Asn Ile Met Lys
35 40 45

Ala Lys Lys Lys Pro Ile Lys Lys Leu Thr Ala Thr Asp Val Gly Val
50 55 60

Asp Leu Ala Pro Arg Gln Gln Val Leu Ser Val Glu Asp Pro Pro Thr 65 70 75 80

Arg Gln Ala Gly Ser Ile Val Pro Asp Val Asp Thr Leu Ile Thr Lys 85 90 95

Leu Lys Glu Lys Gly His Leu 100

<210> 35

<211> 306

<212> DNA

<213> Brassica napus

<220>

<221> CDS

<222> (3)..(305)

<400> 35

gg ttg ggt cga acc ata ggt gga aag ctt ctt tct ctc tcg ctt gac 47
Leu Gly Arg Thr Ile Gly Gly Lys Leu Leu Ser Leu Ser Leu Asp
1 5 10 15

aaa tcc tct ggt tcg ggt ttt cag tcc cat cag gag ttt ctc tat ggt 95 Lys Ser Ser Gly Ser Gly Phe Gln Ser His Gln Glu Phe Leu Tyr Gly 20 25 30

aaa gct gag gtt caa atg aaa ctt gtc cct ggt aac tct gct gga aca 143

<223> GW1

Lys	Ala	Glu	Val 35	Gln	Met	Lys	Leu	Val 40	Pro	Gly	Asn	Ser	Ala 45	Gly	Thr	
_														gag Glu		191
														ctc Leu		239
														gtt Val		287
	_	_	cga Arg		_	С										306
<211 <212)> 36 -> 10 2> PI 3> Bi)1 RT	ica r	napus	5											
)> 36 Gly		Thr	Ile 5	Gly	Gly	Lys	Leu	Leu 10	Ser	Leu	Ser	Leu	Asp 15	Lys	
Ser	Ser	Gly	Ser 20	Gly	Phe	Gln	Ser	His 25	Gln	Glu	Phe	Leu	Tyr 30	Gly	Lys	
Ala	Glu	Val 35	Gln	Met	Lys	Leu	Val 40	Pro	Gly	Asn	Ser	Ala 45	Gly	Thr	Val	
Thr	Thr 50	Phe	Tyr	Leu	Lys	Ser 55	Pro	Gly	Thr	Thr	Trp 60	Asp	Glu	Ile	Asp	
Phe 65	Glu	Phe	Leu	Gly	Asn 70	Ile	Ser	Gly	His	Pro 75	Tyr	Thr	Leu	His	Thr 80	
Asn	Val	Tyr	Thr	Arg 85	Arg	Leu	Trp	Arg	Gln 90	Arg	Thr	Ala	Val	Ser 95	Ser	
Met	Val	Arg	Pro 100	Asp												
<213 <212	0> 3' 1> 2' 2> DI 3> A:	7 NA	ical	Seq	uence	e										
<220	0>															

tgattaatgcctcctctccgttattcg	27
<210> 38	
<211> 27	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> AT3GW2	
<400> 38	
ttgcagtctgagaaattcctccgatcg	27

PCT/GB99/00905

10

15

534 Rec'd PCT/PTO 20 SEP 2000 SIGNAL TRANSDUCTION PROTEIN INVOLVED

IN PLANT DEHISCENCE

This invention relates to novel plant nucleic acid sequences and proteins. The 5 sequences and proteins are useful in the control of plant dehiscence and in the production of male sterile plants.

The production of seed is an important developmental process in all higher plants. In oilseed rape (Brassica napus), following abscission of floral parts, pods or siliques are formed which contain 15-30 seeds. Around 50-70 days after anthesis (DAA) the pods become susceptible to shatter, a process that serves to expel the mature seeds into the surrounding environment. In the days leading to dehiscence, an array of anatomical, molecular and biochemical changes take place, thus preparing both seed and pod for the event. Shatter eventually occurs as a result of a combination of factors including: the creation of tensions within the pod between the lignified valve edge cells of the endocarp and the unlignified dehiscence zone (DZ) cells, weakening of the DZ cell walls by hydrolytic enzyme activity and ultimately due to physical forces such as strong winds or harvesting machinery.

20 Pod development in B. napus can be segmented into three stages. In the first stage, which occurs 0-20 DAA, the newly formed siliques, consisting of two seedcontaining carpels separated by a false septum and a replar region, grow to their full length of around 10cm. The seeds begin to grow when the pods are virtually full size [Hocking and Mason, 1993]. Between 10 and 20 DAA the cells in the replar 25 region begin to differentiate into replar cells, large valve edge cells and form a distinct region, 1-3 cells wide, comprising the DZ [Meakin and Roberts, 1990a].

The second stage occurs between 20 and 50 DAA. From 20 DAA, in conjunction

with termination of pod elongation, secondary cell wall material is deposited in the valve edge cells, and the replar cells become increasingly lignified. The DZ cells do not exhibit thickening of the cell wall. A progressive shrinkage and loss of organelles is apparent in the DZ cells from 40 DAA onwards and eventually these cells separate completely due to hydrolysis of the middle lamella [Meakin and Roberts, 1990a]. In the final stage of pod development, which occurs 50-70 DAA, the lignified cells undergo senescence and the necessary tensions are created so that the desiccated pod, containing mature seed, eventually shatters.

- Molecular studies of the penultimate stage of pod development have revealed a spatial and temporal correlation between the up-regulation of a number of mRNAs and pod dehiscence in B. napus. These mRNAs encode a polygalacturonase (PG) and a proline-rich protein (SAC51). Further analysis of the expression of the PG following fusion of a pod-specific Arabidopsis thaliana PG promoter to GUS
 [Jenkins et al., (1997)], has revealed that reporter gene expression is restricted precisely to the layer of cells comprising the pod DZ in transgenic B. napus. From 40 DAA, Meakin and Roberts (1990b) reported a progressive increase in β-1,4-glucanase (cellulase) activity in the DZ.
- It is understood that the processes of dehiscence and abscission are not regulated by the same environmental or chemical signals, but that they involve controlled degradation of cell wall material and cell separation in a distinct group of cells.

 Both ethylene and indole-3-acetic acid (IAA) appear to be important regulators of the timing of the abscission process but the role of these plant hormones in dehiscence is less clearly defined. The increase in cellulase activity has been shown to correlate with a rise in the production of ethylene, mainly from the seed, which peaks at around 40 DAA [Meakin and Roberts, 1990b; Johnson-Flanagan and Spencer, 1994].

10

15

20

25

Developmental processes, such as pod dehiscence, which involve highly regulated and controlled expression of an array of different genes at a precise time and cellular location, clearly require an intricate signal transduction network.

Further and improved genetic elements to control plant processes in this area are constantly desired. We describe the isolation, for the first time, of a plant cDNA (DZ2) encoding an individual response regulator protein, the expression of which is closely correlated with dehiscence of fruit in *B. napus*. DZ2 has a role in the ability to control molecule signaling during the events leading to shatter and thus to control pod shatter in plants. In addition to the identification of the nucleic acid termed "DZ2" a homologous, but not identical sequence and protein were also identified from *B. napus*. This sequence was designated "DZ2B". Sequence analysis of DZ2 and DZ2B shows that there are two DZ2 genes in *B. napus*, each represented by a slightly different cDNA (here termed DZ2 and DZ2B). This is consistent with one gene being encoded by the *B. campestris* derived genome and the other from the genome derived from *B. oleracea*. In this text, the designation "DZ2" is equivalent to the CW1 designation in UK 9806113.8 (as seen from Figure 1).

According to a first aspect of the invention there is provided nucleic acid optionally encoding a signal transduction protein involved in the process of dehiscence. Such a sequence or signal transduction protein has never previously been described in plant dehiscence.

In this text, the term "involved in the process of dehiscence" means any nucleic acid (preferably) encoding any protein which has an effect in the dehiscence process, in particular a protein or nucleic acid sequence involved in an MAP Kinase cascade or any other protein or nucleic acid sequence which results in changes in the expression of genes involved in dehiscence, such as upregulation of genes encoding

10

15

polygalacturonase, cellulase, senescence-related proteins and/or downregulation of genes encoding for proteins involved in cells wall biosynthesis. The nucleic acid sequences/proteins of the present invention which are "involved in the process of plant dehiscence" are not the individual structural genes or proteins which cause dehiscence (polygalacturonases etc.). Rather, the nucleic acid sequences/proteins of the present invention are sequences/proteins which have an effect on the expression of such structural genes or proteins. One advantage of the present invention is that the use of such nucleic acid sequences/proteins enables the possibility to influence the whole process of dehiscence rather than just one element of it. Preferably the protein or nucleic acid sequence of the present invention which is involved in the process of dehiscence effects a structural protein which is a hydrolytic enzyme such as polygalacturonase or cellulase.

The nucleic acid of the first aspect of the invention may be a nucleic acid which is naturally expressed in a dehiscence zone. Such a nucleic acid will most accurately reflect nucleic acid naturally expressed in a plant. Preferably the dehiscence zone is a pod (also termed "siliques"), anther and/or funiculus dehiscence zone. Preferably the plant is a member of the Brassica family, maize, wheat, soyabean, *Cuphea* or sesame.

20

A second aspect of the invention provides nucleic acid encoding a protein wherein the protein:

a) comprises the amino acid sequence shown in figure 1 or;

25

b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above, but has at least 40% amino acid sequence identical therewith; or

- c) is a fragment of a protein as defined in a) or b) above, which is at least 10 (preferably 20 or 30) amino acids long.
- The percentage amino acid identity can be determined using the default parameters of the GAP computer program, version 6.0 described by Deveraux et al., 1984 and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilises the alignment method of Needleman and Wunsch 1970 as revised by Smith and Waterman 1981. More preferably the protein has at least 45% identity to the amino acid sequence of Figure 1, through 50%, 55% 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% identity using the default parameters.

The skilled person will appreciate that various changes can sometimes be made to the amino acid sequence of a protein (which has a desired property) to produce variants (often known as "muteins") which still have said property. Such variants of the protein describe in a, b and c above are within the scope of the present invention and are discussed in greater detail below in sections (i) to (iii). They include allelic and non-allelic variants.

20 (i) Substitutions

15

An example of a variant of the present invention is a polypeptide as defined in a, b or c above, apart from the substitution of one or more amino acids with one or more other amino acids.

The skilled person is aware that various amino acids have similar characteristics.

One or more such amino acids of a protein can often be substituted by one or more other such amino acids without eliminating a desired property of that protein.

For example, the amino acids glycine, alanine, valine, leucine and isoleucine can often be substituted for one another (amino acids having aliphatic side chains). Of these possible substitutions it is preferred that glycine and alanine are used to substitute for one another (since they have relatively short side chains) and that valine, leucine and isoleucine are used to substitute for one another (since they have larger aliphatic side chains which are hydrophobic). Other amino acids that can often be substituted for one another include phenylalanine, tyrosine and tryptophan (amino acids having aromatic side chains); lysine, arginine and histidine (amino acids having basic side chains); aspartate and glutamate (amino acids having acidic side chains); asparagine and glutamine (amino acids having amide side chains); and cysteine and methionine (amino acids having sulphur containing side chains).

Substitutions of this nature are often referred to as "conservative" or "semi-conservative" amino acid substitutions.

15

20

5

10

(ii) Deletions

Amino acid deletions can be advantageous since the overall length and the molecular weight of a polypeptide can be reduced whilst still retaining a desired property. This can enable the amount of protein required for a particular purpose to be reduced. Proteins according to the present invention, which have such deletion(s) are useful. They may interfere with the normal functioning of DZ2; that is, they may act as dominant negative mutations preventing normal DZ2 functioning and thus be of particular value, for example, in reducing pod shatter.

The amino acid sequence shown in figure 1 has various functional regions. For particular applications of the present invention, one or more of these regions may not be needed and may therefore be deleted.

10

15

20

25

(iii) Insertions

Amino acid insertions relative to a polypeptide as defined in a, b or c above can also be made. This may be done to alter the nature of the protein (e.g. to assist in identification, purification, or expression, as explained below in relation to fusion proteins).

Changes in the protein according to the present invention can produce versions of the protein that are constitutively active. If a protein of the present invention acts on an inhibitor of the release of hydrolytic enzymes, then a constitutively active version would prevent or reduce pod shatter

A protein according to any aspect of the invention may have additional N-terminal and/or C-terminal amino acid sequences. Such sequences can be provided for various reasons. Techniques for providing such sequences are well known in the art. They include using gene-cloning techniques to ligate together nucleic acid molecules encoding polypeptides or parts thereof, followed by expressing a polypeptide encoded by the nucleic acid molecule produced by ligation.

Additional sequences may be provided in order to alter the characteristics of a particular polypeptide. This can be useful in improving expression or regulation of expression in particular expression systems. For example, an additional sequence may provide some protection against proteolytic cleavage. This has been done for the hormone somatostatin by fusing it at its N-terminus to part of the β galactosidase enzyme [Itakwa et al., 105-63 (1977)].

Additional sequences can also be useful in altering the properties of a polypeptide to aid in identification or purification.

For example, a signal sequence may be present to direct the transport of the polypeptide to a particular location within a cell or to export the polypeptide from the cell. Hydrophobic sequences may be provided to anchor a polypeptide in a membrane. Thus the present invention includes within its scope both soluble and membrane-bound polypeptides.

Preferably, the nucleic acid according to the second aspect of the invention encodes a signal transduction protein or a functional portion thereof involved in the process of dehiscence. All preferred features of the first aspect of the invention as described above also apply to the second.

The term protein used in this text means, in general terms, a plurality of amino acid residues joined together by peptide bonds. It is used interchangeably and means the same as polypeptide or peptide.

15

20

25

5

10

The nucleic acid according to the first or second aspect of the invention preferably comprises the sequence set out in figure 1 or a sequence which is 40% or more identical, preferably through 45%, 50%, 55%, 60%, 65%,70%, 75%, 80%, 85%, 90%, 95% to the sequence in Figure 1 at the nucleic acid residue level, using the default parameters of the GAP computer program, version 6.0 described by Deveraux *et al.*, 1984 and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilises the alignment method of Needleman and Wunsch 1970 as revised by Smith and Waterman 1981. Further, the nucleic acid may comprise a fragment of a sequence according to the first or second aspect which is at least 30 bases long also 40, 50, 60, 70, 80, or 90 bases in length. While this nucleic acid is the preferred nucleic acid of the invention, it is well known to those persons skilled in the art that because of the nucleic acid "degenerate code" which encodes nucleic acids, a considerable number of variations in nucleic acid

sequence can be used to encode for proteins according to the first or second aspects of the invention.

The nucleic acid of the first or second aspects of the invention may be isolated or recombinant and may be in substantially pure form. The nucleic acid may be antisense to nucleic acid according to the first or second aspects of the invention. As understood by the person skilled in the art introducing the coding region of a gene in the reverse orientation to that found in nature (antisense) can result in the downregulation of the gene and hence the production of less or none of the gene product. The transcribed antisense DNA is capable of binding to and destroying the function of the sense RNA of the sequence normally found in the cell, thereby disrupting function. Antisense nucleic acid may be constitutively expressed, but is preferably limited to expression in those zones (dehiscence) in which the naturally occurring nucleic acid is expressed.

15

20

25

10

5

The nucleic acid according to the first or second aspects of the invention preferably include a promoter or other regulatory sequence which controls expression of the nucleic acid. Promoters and other regulatory sequences which control expression of a nucleic acid in dehiscence zones are known in the art, for example described in WO96/30529 and WO94/23043. Further promoters or other regulatory sequences can be identified and can also include the promoter or other regulatory sequence which controls expression of a nucleic acid as set out in figure 1. The person skilled in the art will know that it may not be necessary to utilize the whole promoter or other regulatory sequence. Only the minimum essential regulatory elements may be required and in fact such elements can be used to construct chimeric sequences or promoters. The essential requirement is, of course, to retain the tissue and/or temporal specificity.

10

15

20

25

The nucleic acid according to the first or second aspects of the invention may be in the form of a vector. The vector may be a plasmid, cosmid or phage. Vectors frequently include one or more expressed markers which enable selection of cells transfected (or transformed: the terms are used interchangeably in this text) with them and preferably, to enable a selection of cells containing vectors incorporating heterologous DNA. A suitable start and stop signal will generally be present and if the vector is intended for expression, sufficient regulatory sequences to drive expression will be present. Nucleic acid according to the first and second aspects of the invention is preferably for expression in plant cells and thus microbial host expression is perhaps less important although not ruled out. Microbial host expression and vectors not including regulatory sequences are useful as cloning vectors.

A third aspect of the invention relates to a cell comprising nucleic acid according to the first or second aspects of the invention. The cell may be termed as "a host" which is useful for manipulation of the nucleic acid, including cloning.

Alternatively, the cell may be a cell in which to obtain expression of the nucleic acid, most preferably a plant cell. The nucleic acid can be incorporated by standard techniques known in the art in to cells. Preferably nucleic acid is transformed in to plant cells using a disarmed Ti plasmid vector and carried by an Agrobacterium by procedures known in the art, for example as described in EP-A-0116718 and EP-A-0270822. Foreign nucleic acid can alternatively be introduced directly into plant cells using an electrical discharged apparatus or by any other method that provides for the stable incorporation of the nucleic acid into the cell. Preferably the stable incorporation of the nucleic acid is within the nucleic DNA of any cell preferably a plant cell. Nucleic acid according to the first and second aspects of the invention preferably contains a second "marker" gene that enables identification of the nucleic acid. This is most commonly used to distinguish the transformed plant cell

containing the foreign nucleic acid from other plants cells that do not contain the foreign nucleic acid. Examples of such marker genes include antibiotic resistance, herbicide resistance and Glucuronidase (GUS) expression. Expression of the marker gene is preferably controlled by a second promoter which allows expression of the marker gene in cells other than those than dehiscence zones (if this is the tissue specific expression of the nucleic acid according to the first or second aspects of the invention). Preferably the cell is from any of the Brassica family (most preferably B. napus), maize, wheat, soyabean, Cuphea and sesame.

- A third aspect of the invention includes a process for obtaining a cell comprising nucleic acid according to the first or second aspects of the invention. The process involves introducing said nucleic acid into a suitable cell and optionally growing on or culturing said cell.
- 15 A fourth aspect of the invention provides a plant or a part thereof comprising a cell according to the third aspect of the invention. A whole plant can be regenerated from the single transformed plant cell by procedures well known in the art. The invention also provides for propagating material or a seed comprising a cell according to the third aspect of the invention. The invention also relates to any plant or part thereof including propagating material or a seed derived from any aspect of the invention. The fourth aspect of the invention also includes a process for obtaining a plant or plant part (including propagating material or seed, the process comprising obtaining a cell according to the third aspect of the invention or, indeed, plant material according to the fourth aspect of the invention and growth (to the required plant, plant part, propagating material etc). Techniques for such a process are commonplace in the art.

A fifth aspect of the invention provides a signal transduction protein involved in the

process of the plant dehiscence. The signal transduction protein according to the fifth aspect may have one or more of the preferred features according to the first or second aspects of the invention. Preferably it may be isolated, recombinant or in substantially pure form. It may comprise the various changes according to the first or second aspects. Preferably the protein is expressed from nucleic acid according to the first or second aspects. Alternatively, the protein can be provided using suitable techniques known in the art.

A sixth aspect of the invention provides a protein which;

a) comprises the amino acid sequence shown in figure 1 or;

b) has one or more amino acid deletions, insertions, or substitutions relative to a protein as defined in a) above and has at least 40% amino acid sequence identity therewith;

15

20

10

5

or a fragment of a protein as defined in a) or b) above which is at least 10 amino acids long. The percentage amino acid identity can be determined using the default parameters of the GAP computer program, version 6.0 described by Deveraux et al., 1984 and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilises the alignment method of Needleman and Wunsch 1970 as revised by Smith and Waterman 1981. More preferably the protein has at least 45% identity to the amino acid sequence of Figure 1, through 50%, 55% 60%, 65%, 70%, 75% 80%, 85%, 90%, 95% identity using the default parameters.

The protein is preferably a signal transduction protein involved in the process of plant dehiscence and again, the preferred features of aspects one, two and five also applied to the sixth aspect.

10

15

20

25

The seventh aspect of the invention provides a process for regulating/controlling dehiscence in plant or in a part thereof, the process comprising obtaining a plant or a part thereof according to the fourth aspect of the invention. The process of dehiscence can be regulated and/or controlled by increasing or decreasing the expression of nucleic acid sequences according to the first or second aspect of the invention. Increased or decreased expression can easily be influenced by the person skilled in the art using technology well known. This includes increasing the numbers of copies of nucleic acid according to the invention in a plant or a plant thereof or increasing expression levels of copies of the nucleic acid present in particular parts or zones of the plant. Preferably the zones are dehiscence zones.

The process according to the seventh aspect of the invention includes obtaining a plant cell according to the third aspect of the invention or part of a plant according to the fourth aspect in the invention and deriving a plant therefrom. Alternatively, the process may comprise obtaining propagating material or a seed according to the fourth aspect of the invention and deriving a plant therefrom.

Preferably, the process of the seventh aspect of the invention is in the pod or the anther of a plant. All preferred features of aspects one to six also apply to the seventh.

An eighth aspect of the invention provides for the use of nucleic acid according to the first to seventh aspects of the invention in the regulation/control of plant dehiscence. All preferred features of aspects one to seven also applies to the eighth.

The ninth aspect of the invention provides for the use of nucleic acid according to the first or second aspect of the invention as a probe. Such a probe can be used in techniques well known in the art to identify the presence of identical or homologous

nucleic acid sequences from any source, preferably a plant source. The ninth aspect of the invention also provides nucleic acid identified by use of the nucleic acid from aspects one or two as a probe.

- A tenth aspect of the invention provides for the use of nucleic acid according to aspects one or two of the invention in the production of a cell, tissue, plant or part thereof, or propagating material. Again, all preferred features of aspects one and two also apply to the tenth.
- An eleventh aspect of the invention provides for nucleic acid comprising one or more of the underlined sequences as set out in figure 1 or the primer sequences in Fig 5, Fig 9 or Fig 11. Such nucleic acid sequences are preferably used as primers in an PCR (Polymerase Chain Reaction) process in order to amplify nucleic acid sequences.

A twelfth aspect of the invention provides the use of nucleic acid according to the first or second aspects of the invention to identify another other protein or proteins which interact with its expression product. Such use can be carried out by the yeast two hybrid screening method (or others known in the art). The yeast two hybrid screening method is described for this aspect of the invention, in general, with reference to the sequence described as DZ2. A potential way to implement the yeast 2-hybrid screen is outlined, as follows:

DZ2 is linked to the Gal4 DNA binding domain and expressed in yeast which contains a pGAL4-lacZ gene. For activity of lacZ a second protein is required that contains the DNA transcriptional activation domain of GAL4 and that interacts with the DZ2 protein. This is provided by making a cDNA expression library from plant DZ zones which results in fusions of plant

proteins to the GAL4 activation domain. This library is transformed into the yeast strain that contains pGAL4-LacZ and expresses the DZ2-Gal4 DNA bining domain protein fusion. Colonies that have lacZ activity are transformed with a gene for a protein that interacts with DZ2.

5

Using such a system, upstream and downstream components of any signal transduction pathway can be identified, thus resulting in further ability to control/regulate dehiscence and/or male sterility.

A thirteenth aspect of the invention provides for a protein, as defined according to the limitations of the second aspect of the invention (without reference to figure 1) and nucleic acid encoding the protein, wherein the protein is capable of being identified according to the use (or method) according to the twelfth aspect of the invention.

15

20

25

A fourteenth aspect of the invention provides for the use of a protein according to the fifth or sixth aspect of the invention as a probe. In this context the probe is a means to identifying interacting entities (such as other proteins), including upstream and downstream interacting signal components. A protein according to the fifth or sixth aspect of the invention can be used as a probe to directly look for interactions with other proteins, i.e. purified protein can be used to look for complex formation with other plant protein, particularly isolated from the DZ zone. For example, a modified recombinant DZ2 protein can be made with a sequence tag, such as a Histag, that enables the DZ2 + interacting protein to be directly purified on a His affinity column. Alternatively, an antibody can be raised to DZ2 protein. This antibody is then used to identify DZ2 protein complexes and to purify the complexes. The DZ2 interacting proteins can be purified and microsequenced to enable cloning of the genes for these interacting proteins.

10

15

20

25

The present invention provides a particularly useful method by which plant dehiscence can be regulated/controlled.

In addition to the use of the present invention in the production of shatter resistance or shatter-delayed plants such as oil seed rape, the invention may be used to control/regulate pollen release (by the control/regulation of anther dehiscence) which can produce male sterile plants. The temporal and spatial expression of nucleic acid encoding a protein according to the first and second aspects of the invention may require adjustment in obtaining the correct levels of dehiscence delay or prevention in different zones. For example, if pod dehiscence is required but anther dehiscence is not, it is necessary to ensure that expression of nucleic acid according to a first or second aspect of the invention has the correct temporal and spatial expression in order to obtain pod dehiscence or delay but not, to any substantial extent, anther dehiscence. This can be obtained by processes known in the art and may require use of particular promoter sequences to obtain the desired result. Usually in plant transformation, some difference in the level of expression of nucleic acid is observed in different plants. In some cases, the ratio of expression levels in different tissues can vary between different plant transformants thus providing essentially tissuespecific expression in one or other of the target tissues in some of the plant transformants. In the present invention, the natural expression of nucleic acid according to the first or second aspects may be predominantly higher in pod dehiscence zones and lower in the anther and funiculus dehiscence zones. However, as described above, it is possible to obtain plants in which the protein expression is regulated in a particular dehiscence zone. Accordingly, a particularly useful aspect of the invention is the provision of plants which have one or both of the following features; are male sterile, are shatter resistant.

As described earlier, the process of dehiscence at the dehiscence zone involves the secretion of a number of enzymes, including hydrolytic enzymes. While previous attempts have been made to down or up regulate specific genes encoding particular proteins involved in the process of dehiscence, regulation by means of a signal transduction protein which effects expression of a number of genes is likely to be more effective than regulation of a single gene. In addition to this, the nucleic acid of the present invention has been identified as being expressed earlier than several other known genes involved in the process of plant dehiscence. This suggests that it is important earlier on in the process of plant dehiscence and can be used to control/regulate plant dehiscence at an earlier stage.

The nucleic acid encoding a signal transduction protein involved in the process of dehiscence or the signal transduction protein itself may be a component of a signal pathway that may either positively or negatively regulate pod shatter.

15

10

5

A more detailed explanation of such regulations/control, described with reference to a pod shatter (dehiscence) model is described below. As a skilled person will acknowledge, the model described below also relates to other general processes of dehiscence such as in the anther.

20

25

In the process of dehiscence, a particular signal transduction protein may be required to transmit a signal from the almost mature seed which initiates the expression or release of enzymes required for pod shatter. In this model, developmental signals switch on expression and/or activation of a particular signal transduction protein in the pod dehiscence zone. This leads to expression of genes required for the release of pod dehiscence zone enzymes (such as hydrolytic enzymes). In this case, prevention of activity of the signal transduction protein, for example by downregulation of expression of this protein, would result in reduced dehiscence.

15

20

25

Alternatively, the developing seed may transmit a signal which represses the expression and/or activity of a particular signal transduction protein until late in cell development. In this model, developmental signals switch on a particular signal transduction protein which, in due course, represses the expression of genes required for release of dehiscence zone specific enzymes (such as hydrolytic enzymes). In this case, expression of a modified signal transduction protein that is constitutively active would result in reduced dehiscence.

A signal transduction protein which is either positively or negatively involved in the process of dehiscence can be used according to the present invention.

In addition to DZ2 several other DZ-expressed genes have been previously isolated and individually downregulated to result in B. napus plants that have increased resistance to pod shatter; namely Sac66 (WO 96/30529 - Figure 15), DZ15 (Figure 16) and OSR 7(9) (Figure 17). It is anticipated that downregulation of more than one gene involved in pod shatter will further increase resistance to pod shatter. This could be achieved by combining different transgenes by transformation with several transgenes each designed to downregulate a different DZ-expressed gene or by crossing together B. napus lines that individually are transformed with such transgenes. Such methods are complex either involving transformation with a construct containing multiple chimeric genes or require the maintenance of several transgenic loci in the breeding program. A preferred method is to transform with a chimeric gene consisting of a single promoter driving expression of an antisense or partial sense transcript which is comprised of elements of all the DZ-expressed genes to downregulated. Similarly a single promoter could be used to drive the expression of multiple ribozymes each targeted against a different DZ-expressed gene. The use of a single promoter to drive expression of a combination of antisense, partial sense

and ribozymes is also possible. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. A suitable DZ-specific promoter would be that of DZ2, DZ2B, DZ2AT3 or ESJ2A (WO 99/13089).

5

10

15

Accordingly, the present invention provides a nucleic acid sequence according to the first or second aspects (and also all aspects which include the first or second aspects) in combination with one or more further nucleic acid sequences which are dehiscence-zone expressed. Examples of such sequences include Sac66, DZ15 and OSR(7), Figures 15-17 respectively. Such sequence may be in sense or in antisense orientation. Such a sequence may be included as full length genomic, full-length cDNA or partial sequences; the sequences may be as shown in the figures or may have the same sequence identity (both for aminoacid sequence and nucleic acid sequence) as described above for the protein according to the second aspect of the invention or the nucleic acid according to the first or second aspects of the invention. As will be recognised by those skilled in the art a partial sequence may be useful in either the sense or antisense orientation.

The invention is described by reference to the enclosed drawings;

20

- Figure 1 DZ2 full length sequence showing original PCR product and primer sites
- Figure 2 Amino acid alignment with bacterial response regulator proteins & ETR1

	Figure 3	Northern analysis of expression of DZ2 in pods and other tissues.
		The lower panel shows the ethidium bromide-stained RNA gel prior to blotting and probing with DZ2
5	Figure 4	Comparison of bacterial two-component systems with DZ2
	Figure 5	Sequence of the promoter region of B.napus DZ2B.
10	Figure 6	Nucleic and putative peptide sequence alignments of DZ2 with DZ2B.
	Figure 7	Northern analysis of expression of DZ2B in pods and other tissues. The probe was labeled DZ2B cDNA.
15	Figure 8	Schematic diagram of pDZ2B-GUS-SCV
	Figure 9	DZ2AT3 cDNA sequence showing the putative DZ2AT3 peptide.
•	Figure 10	Amino acid alignment of DZ2AT3 with DZ2 and DZ2B.
20	Figure 11	Sequence of the promoter region of A.thaliana DZ2AT3.
	Figure 12	Schematic diagram of pDZ2AT3-GUS-SCV.
25	Figure 13	Schematic diagram of pPGL-DZ2as-SCV and pDZ2B-DZ2as-SCV.
	Figure 14	Schematic diagram of pWP357-SCV.
	Table 1	Pod shatter resistance of WP357-SCV transformants.

10

- Figure 15 Nucleic acid sequence and putative amino acid sequence of Sac66.
- Figure 16 Nucleic acid sequence and putative amino acid sequence of DZ15.

Figure 17 Nucleic acid sequence and putative amino acid sequence of OSR7 (9)

The present invention is now described with reference to the following, non-limiting examples.

Example 1- Isolation and characterisation of expression of DZ2

Plant Material

Seeds of *B. napus* cv Rafal were grown as described by Meakin and Roberts, (1990a) with the following modifications. Single seedlings were potted into 10cm pots, and after vernalization, were re-potted into 21cm pots. At anthesis tags were applied daily to record flower opening. This procedure facilitated accurate age determination of each pod. Pods were harvested at various days after anthesis (DAA). The dehiscence zone was excised from the non-zone material and seed using a scalpel blade (Meakin and Roberts (1990b)) and immediately frozen in liquid N₂ and stored at -70°C.

RNA Isolation

All chemicals were molecular biology grade and bought from either Sigma Chemical Ltd (Dorset, UK), or Fisons (Loughborough, UK). Total RNA was extracted using the polysomal extraction method of Christoffersen and Laties, (1982), with the following alterations. The plant material was ground to a powder in liquid N₂ and then in 10 volumes of extraction buffer (200mM Tris-acetate [pH 8.2], 200mM magnesium

acetate, 20mM potassium acetate, 20mM EDTA, 5% w/v sucrose, after sterilisation 2mercaptoethanol was added to 15mM and cycloheximide added to a final concentration of 0.1 mg ml⁻¹). The supernatant was then layered over 8 ml 1M sucrose made with extraction buffer and centrifuged in a Kontron™ (Switzerland) TFT 70.38 rotor at 45,000rpm (150,000g) for 2 hr at 2°C in a Kontron CENTRIKON™ T-1065 ultra-5 centrifuge. Pellets were then resuspended in 500 µl 0.1M sodium acetate, 0.1% SDS, pH 6.0 and phenol/chloroform (1:1 v/v) extracted and the total RNA precipitated. Poly(A)+ RNA was isolated from total RNA extracted, from both the zone and nonzone tissue of 40, 45 and 50 DAA pods, using a Poly(A) QUIK[™] mRNA purification kit (Stratagene, Cambridge, UK) following the manufacturers instructions, and then bulked together. Total RNA was also extracted from leaves, stems, seeds and pods using a method described by Dean et al, (1985) for use in Northern analyses.

Differential display

15

20

25

10

This was performed essentially as described by Liang and Pardee (1992) using RNA extracted from 40 DAA pod dehiscence zones and non-zones. First strand cDNA copies of the RNAs (40 DAA DZ/NZ) were made using 50U M-MLV (Moloney Murine Leukemia Virus) reverse transcriptase (50U/μL) (Stratagene) in a 20μL reaction containing 1x M-MLV buffer, 2.5mM dNTPs (Pharmacia), 1µg RNA, 30U RNAse inhibitor (Promega) and 10μM oligo dT anchor primer 7 (5'-TTTTTTTTTTTGG-3'). The reaction conditions were as follows: 65°C for 5 minutes, 37°C for 90 minutes and 95°C for 5 minutes. Following first strand cDNA synthesis, $60\mu L$ dH2O were added and the samples were either used directly for PCR or stored at -20°C.

For PCR, 2µL cDNA were used as template in a 20µL reaction containing 1x PCR buffer, 1mM MgCl₂, 2µM dNTPs, 10µM oligo dT anchor primer 7 (5'-

TTTTTTTTTTTGG-3'), 2.5µM arbitrary primer A (5'- AGC CAG CGA A -3'). 0.5μL 35S-dATP (>1000 Ci/mmoi) (Amersham) and 1U Tag DNA polymerase $(5U/\mu L)$ (Gibco BRL). The thermocycling conditions were as follows: 40 cycles of 94°C for 30 seconds, 40°C for 2 minutes, 72°C for 30 seconds followed by 72°C for 5 5 minutes. The PCR products were fractionated on a 5% polyacrylamide/7M urea gel after addition of 5µL loading buffer (95% (v/v) formamide, 20mM EDTA. 0.05% (w/v) xylene cyanol, 0.05% (w/v) bromophenol blue) to each sample. Following electrophoresis the gel was dried at 80°C under vacuum for 1 hour then exposed to X-ray film (BioMax-MR, Kodak) in a light tight cassette for 48 hours. 10 The dried gel and autoradiogram were aligned so that bands that appeared in the DZ and not in NZ could be cut out and the DNA eluted according to Liang et al. (1995). The eluted PCR products $(4\mu L)$ were reamplified in a $40\mu L$ reaction containing 1x PCR buffer, 1mM MgCl₂, 20µM dNTPs, 10µM oligo dT anchor primer 7 (5'-TTTTTTTTTTTGG-3'), 2.5µM arbitrary primer A (5'- AGC CAG CGA A -3') 15 and 2U Taq DNA polymerase (5U/µL) (Gibco BRL) using the following thermocycling conditions: 40 cycles of 94°C for 30 seconds, 40°C for 2 minutes. 72°C for 30 seconds followed by 72°C for 5 minutes. The resulting PCR product was cloned into the TA cloning vector (Invitrogen) and sequenced (Figure 1). In order to prepare an antisense strand-specific riboprobe, the PCR product was 20 subcloned into pBluescript (Stratagene).

Expression analysis and characterisation of DZ2

Northern analysis using an antisense strand-specific riboprobe to the DZ2 PCR product, showed that DZ2 hybridised to a transcript of 0.6kb which is expressed in the DZ of 20-50 DAA pods with a peak in expression at 40DAA. Minimal expression was observed in the pod NZ [Figure 2]. A random-primed labelled DNA probe (Stratagene) of the 330bp DZ2 PCR product (amplified using primers DZ2FL

10

15

20

25

and DZ2RL - see Figure 1) was used to screen a *B. napus* DZ cDNA library from which, following three rounds of screening to obtain pure plaques, a full length DZ2 cDNA (606bp) was obtained (Figure 1). An antisense strand-specific riboprobe of the full length DZ2 cDNA was hybridised to total RNA extracted from pod DZ/NZ (as in Figure 2), leaf abscission zones (AZ) and non-zones (NZ) (following exposure to 10μ L/L ethylene for 72 hours), seed, root, flower and leaf. Figure 3 shows that DZ2 hybridises to a 0.6kb message which is present in the pod DZ at 20-50 DAA with maximum expression at 40DAA. Again there is minimal expression in pod NZ and no apparent expression of DZ2 in AZ, NZ, leaf, root, seed or flower RNA. By the sensitive technique of RT-PCR analysis DZ2 expression can also be detected in anthers and the funiculus, both tissues that contain dehiscent zones

The 606bp cDNA (DZ2) encodes a putative protein of 136 amino acids. Comparison of the DZ2 translated sequence to the OWL protein database [Bleasby and Attwood (1994)] showed low but consistent homology to a group of bacterial proteins comprising two-component regulatory systems. In particular, DZ2 possesses the conserved amino acid residues required for phosphorylation of the receiver domain of the response regulator component (see Figure 4). DZ2 plays a role in a signal transduction cascade resulting at least in one respect in pod shatter. It is therefore a good candidate for down-regulation of pod shatter processes using antisense technology. DZ2 is a novel plant protein in that independent proteins with homology to bacterial receivers are yet to be reported in plants.

The full length cDNA was excised from the pBluescript cloning vector by digestion with *Eco*RI and *Xho*I restriction enzymes (Gibco BRL). Following purification from a 1% agarose gel the 606bp cDNA was random primed labelled (Stratagene) and used to screen a *B. napus* genomic library in the BlueStar vector. Following three rounds of screening to obtain pure plaques, a single genomic clone was isolated

10

15

which carries a 15kb genomic DNA insert. The promoter of the DZ2 gene is isolated from this genomic clone using standard techniques (see Example 2).

Example 2 - Isolation and characterisation of the B.napus DZ2B promoter.

To obtain the B.napus DZ2 promoter a B.napus genomic library was screened with a labelled DZ2 probe. The full length cDNA was excised from the pBluescript cloning vector (Stratagene) by digestion with *EcoRI* and *XhoI* restriction enzymes (Gibco BRL). Following purification from a 1% agarose gel the 606bp cDNA was random primed labelled (Stratagene) and used to screen a *B. napus* genomic library in the BlueStar vector (Novagen). Following three rounds of screening to obtain pure plaques, a single genomic clone was isolated which carries a 15kb genomic DNA insert. The region hybridising to DZ2 was sequenced and found to encode a protein homologous to, but not identical to DZ2. This DZ2-like gene was designated DZ2B (Figure 5). The primers DZ2BFL (Figure 5) and T7 were used to PCR out a DZ2B cDNA from the B.napus DZ cDNA library.

- 5' AACCAAGTCAGTAGAAGTG 3' DZ2BFL
- 5' AATACGACTCACTATAGG 3' T7

20

25

The DZ2 and DZ2B cDNAs are 80% identical (according to the default parameters of the GAP computer program, version 6, Deveraux et al., 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) at the nucleotide level in the region of overlap of the coding sequences (Figure 6a) and the putative proteins encoded by DZ2 and DZ2B are 80% identical (according to the default parameters of the GAP computer program, version 6, Deveraux et al., 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) (Figure 6b). Sequence analysis of more DZ2 and DZ2-like cDNAs and Southern analysis shows that there are two DZ2 genes in B.napus, DZ2 and DZ2B,

each represented by 2 slightly different cDNAs. This is consistent with one gene being encoded by the B.campestris derived-genome and the other from the genome derived from B.oleracea.

- RT-PCR with primers specific to DZ2B showed that DZ2B is only expressed in pods. This was confirmed by northern analysis which showed preferential expression in the DZ (Figure 7). Thus DZ2B has a similar pattern of expression as DZ2 and is thus a suitable source of a DZ-expressed promoter.
- Primers DZ2BGenF and DZ2BGenR were used to PCR a 1253bp DZ2B promoter fragment (Figure 5).
 - 5' GGCTCTAGACGAACTGCGGAGCAAGG 3' DZ2BGENF
 - 5' CTGCCATGGTCGGTTTTTTTTTCTTCGAAC 3' DZ2BGENR

15

20

These primers introduced an XbaI site at the 5' end of the PCR fragment and an NcoI site around the initiating Met of DZ2B. Thus the PCR fragment was cloned as an XbaI, NcoI fragment between the XbaI and NcoI sites of pWP272 (WO 99/10389) forming pDZ2B-GUS. The chimeric pDZ2B-GUS-CaMV polyA gene was then transferred as an XbaI, XhoI fragment between the XbaI and SalI sites of pSCV nos-nptII (WO 95/20668) forming pDZ2B-GUS-SCV (Figure 8). The pDZ2B-GUS-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed B.napus plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989). Gus expression is observed in the pod DZ.

25

10

15

20

25

Example 3 - Isolation and characterisation of a DZ2 Arabidopsis thaliana homologue

To demonstrate that a DZ2 orthologous gene can be isolated from another plant species the functional equivalent of B.napus DZ2 / DZ2B was isolated from Arabidopsis thaliana. The B.napus DZ2 cDNA was used as a probe to screen an Arabidopsis cDNA library (J. Giraudat, ISV-CNRS, France). Figure 9 shows the sequence of a cDNA (DZ2AT3) that hybridised to the DZ2 probe. DZ2AT3 has 85% nucleic acid identity to DZ2 and 85% to DZ2B (according to the default parameters of the GAP computer program, version 6, Deveraux et al., 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) in the coding regions which are common to all three sequences. The putative peptide encoded by DZ2AT3 has 80% identity to DZ2 and 80% to DZ2B (according to the default parameters of the GAP computer program, version 6, Deveraux et al., 1984, and available from the University of Winsconsin Genetics Computer Group (UWGCG)) in the regions which are common to all three sequences (Figure 10). RT-PCR analysis of RNA isolated from leaves, roots, flowers and siliques showed that DZ2AT3 was specifically expressed in siliques. Southern hybridisation analysis showed that the DZ2AT3, DZ2 and DZ2B probes each identify a single identical band in A.thaliana. This indicates that A.thaliana contains one DZ2 gene in contrast to B.napus which contains two.

The Genome walker kit (Clonetech) was used to isolate the DZ2AT3 promoter from A.thaliana genomic DNA. Nested PCR was performed using primer GW1 first, then AT3GW2 each in conjunction with the Genome Walker kit primer (Figure 9). Figure 11 shows the sequence of the promoter region of DZ2AT3 thus obtained. The primers ATDZ2F and ATDZ2R were used to PCR a 1195bp promoter fragment from the DZ2AT3 genomic sequence (Figure 11).

10

15

20

25

5' CACTAGTAGGGCACGCGTGGTCG 3' ATDZ2F

5' TCCATGGTCGATTTCTTTTCTCTCAAG 3' ATDZ2R

These primers introduced an SpeI site at the 5' end of the PCR fragment and an NcoI site around the initiating Met of DZ2AT3. Thus the PCR fragment was cloned as an SpeI, NcoI fragment between the XbaI and NcoI sites of pWP272 (WO 99/13089) forming pDZ2AT3-GUS. The chimeric pDZ2AT3-GUS-CaMV polyA gene was then transferred as a Sal1, XhoI fragment into the SalI site of pSCV nosnptII (WO 95/20668) forming pDZ2AT3-GUS-SCV (Figure 12). The pDZ2AT3-GUS-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed B.napus plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989). Gus expression is observed in the pod DZ.

Example 4 - Production of shatter-resistant *B.napus* plants by antisense downregulation of DZ2

Downregulation of the DZ2 gene or reduction in DZ2 protein levels in the pod DZ will result in plants that are resistant (or more resistant than without this modification) to pod shatter. Standard techniques, commonplace in the art, such as the expression of antisense DZ2 mRNA, full sense mRNA, partial sense mRNA or a ribozyme directed against DZ2 mRNA are effective. Expression of these RNAs requires a promoter that is active in the pod DZ at the time at which DZ2 is expressed. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. A suitable promoter would be that of DZ2. Although DZ2 is expressed in the anther DZ, pod DZ and funiculus DZ, DZ2 promoter -GUS fusion studies show that in different transformants the relative level of expression in these three sites is variable but is stability hereditable. Thus some transformants are obtained in which expression is largely or exclusively

10

confined to the pod DZ. This suggests that the pDZ2 promoter is comprised of distinct elements each specifying expression in a particular DZ. Alternatively the site of transgene integration may influence relative expression levels in the DZ tissues. The DZ2 promoter is therefore linked to the DZ2 cDNA such that the DZ2 is in the antisense orientation forming pDZ2-antiDZ2. This chimeric gene is transferred to the binary vector pNos-NptII-SCV (W0 96/30529). This binary vector is transferred to the agrobacterial strain pGV2260 and transformed *B.napus* plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989) Plant Cell Reports 8, 238-242. A proportion of transformed *B.napus* plants exhibit reduced levels of DZ2 message and are resistant to pod shatter.

Example 5 - Production of shatter-resistant B.napus plants by antisense downregulation of DZ2

15 Downregulation of the DZ2 gene or reduction in DZ2 protein levels in the pod DZ will result in plants that are resistant to pod shatter. Techniques such as the expression of antisense DZ2 mRNA, full sense mRNA, partial sense mRNA or a ribozyme directed against DZ2 mRNA will be effective. Expression of these RNAs requires a promoter that is active in the pod DZ at the time at which DZ2 is 20 expressed. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. Although DZ2 / DZ2B is expressed in the anther DZ, pod DZ and funiculus DZ, DZ2B promoter -GUS fusion studies show that in different transformants the relative level of expression in these three sites is variable but is stably hereditable. Thus some transformants are obtained 25 in which expression is largely or exclusively confined to the pod DZ. This suggests that the pDZ2 promoter is comprised of distinct elements each specifying expression in a particular DZ. Alternatively the site of transgene integration may influence relative expression levels in the DZ tissues. Thus a suitable DZ-specific promoter would be that of DZ2, DZ2B, DZ2AT3 or ESJ2A (WO 99/13089).

The primers DZ2FLA and DZ2RLA were used to PCR a 349bp fragment from the DZ2 cDNA:-

5 5' GGCGAATTCCGGTGAGGAGGCAGTAATC 3'

DZ2FLA

5' GGCCCATGGCATACATACACACTTAGAC 3'

DZ2RLA

The primers introduce an EcoRI and NcoI site at the ends of the DZ2 PCR fragment.

To link the DZ2 PCR fragment in an antisense orientation to the promoter of ESJ2A

(pPGL) the DZ2 PCR fragment was cloned as a NcoI, EcoRI fragment between the NcoI and EcoRI sites of pWP272 (WO 99/13089) forming pPGL-DZ2as. The pPGL-antisense DZ2 chimeric gene was transferred as a XbaI, XhoI fragment from pDZ2as into the XbaI and SaII sites of the binary vector pSCV nos-nptII (WO 95/20668) forming pPGL-DZ2as-SCV (Figure 13a).

15

20

25

The pPGL-DZ2as-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed B.napus plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989). A proportion of transformed B.napus plants exhibit reduced levels of DZ2 and DZ2B message and were resistant to pod shatter

Similarly, to link the DZ2 PCR fragment in an antisense orientation to the promoter of DZ2B, the DZ2 PCR fragment is cloned as a NcoI, EcoRI fragment between the NcoI and EcoRI sites of pDZ2B-GUS forming pDZ2B-DZ2as. The pDZ2B-DZ2as chimeric gene is transferred as a XbaI, XhoI fragment from pDZ2B-DZ2as into the XbaI and SaII sites of the binary vector pSCV nos-nptII (WO 95/20668) forming pDZ2B-DZ2as-SCV (Figure 13b).

The pDZ2B-DZ2as-SCV binary vector is transferred to the agrobacterial strain pGV2260 and transformed B.napus plants. Again a proportion of transformed B.napus plants exhibit reduced levels of DZ2 and DZ2B message and are resistant to pod shatter.

5

15

20

25

Similarly a proportion of B.napus plants transformed with a pDZ23A-DZ2as-SCV construct exhibit reduced levels of DZ2 and DZ2B message and are resistant to pod shatter.

10 Example 6 - Production of shatter-resistant B.napus plants by antisense downregulation of multiple DZ-expressed genes

In addition to DZ2 several other DZ-expressed genes have been previously isolated and individually downregulated to result in B.napus plants that have increased resistance to pod shatter; namely Sac66 (WO 96/30529 Figure 15), DZ15 (Figure 16) and OSR 7(9) (Figure 17). It is anticipated that downregulation of more than one gene involved in pod shatter will further increase resistance to pod shatter. This could be achieved by combining different transgenes by transformation with several transgenes each designed to downregulate a different DZ-expressed gene or by crossing together B.napus lines that individually are transformed with such transgenes. Such methods are complex either involving transformation with a construct containing multiple chimeric genes or require the maintenance of several transgenic loci in the breeding program. A preferred method is to transform with a chimeric gene consisting of a single promoter driving expression of an antisense or partial sense transcript which is comprised of elements of all the DZ-expressed genes to be downregulated. Similarly a single promoter could be used to drive the expression of multiple ribozymes each targeted against a different DZ-expressed gene. The use of a single promoter to drive expression of a combination of

antisense, partial sense and ribozymes is also possible. Ideally the promoter will be pod DZ-specific, however a useful promoter may be pod-specific or even constitutively active. A suitable DZ-specific promoter would be that of DZ2, DZ2B, DZ2AT3 or ESJ2A.

5

10

Consequently the ESJ2A promoter was linked to a multiple antisense gene consisting of elements of Sac66, DZ2, DZ15 and OSR 7(9) in the following manner:- The original DZ15 PCR product in pCRII (Invitrogen) (see Figure 16) was cloned as an EcoRI fragment into pBluescript SK (Stratagene) forming pDZ15-BS, such that the DZ15 3' end is nearest the SstI site of the vector . T7 and DZ15RL primers were used to PCR a 456bp DZ15 fragment from pDZ15-BS which was cloned into the EcoRV site of pGEM5zf (Promega) forming pWP351, such that the DZ15 3' end is nearest the SphI site of the vector.

- 15 5' AATACGACTCACTATAGG 3' T7
 - 5' AACAGCTGAAAACCTCACGAAG 3' DZ15RL

The EcoRI, NcoI fragment of pWP351 cloned between the EcoRI and NcoI sites of pDZ2-BS forming pWP356. pDZ2-BS consists of the DZ2 cDNA cloned as an EcoRI, XhoI fragment into pBluescript SK such that the 3' end is nearest the KpnI site of the vector. A 361bp Sac66 fragment was PCRed from the Sac66 cDNA (WO 96/30529) using the primers F1 and RI which introduce NcoI and PstI sites into the ends of the PCR product.

- 25 5' GGCCCATGGCTGCCAAGCTTTGAGTAGC 3' F1
 - 5' GGCCTGCAGTGCCTAGGATCTGGAAGC 3' RI

10

The Sac66 PCR product was cloned as an NcoI, EcoRI fragment between the NcoI and EcoRI sites of pWP272 (WO 99/13089) forming pWP288A. EcoRI DZ15+DZ2 and OSR 7(9) fragments from pWP356 and pOSR 7(9)-CRII were cloned into the EcoRI site of pWP288A such that DZ15+DZ2 and OSR 7(9) are in an antisense orientation with respect to PGL promoter. (pOSR 7(9)-CRII consists of the 306bp OSR 7(9) PCR fragment (see Figure 17) cloned into pCRII (Invitrogen)). The chimeric pPGL-antisense Sac66+DZ2+Dz15+OSR 7(9) gene was transferred as a XbaI. XhoI fragment into the XbaI and SalI sites of the binary vector pSCV nosnptII (WO 95/20668) forming pWP357-SCV (Figure 14). The pWP357-SCV binary vector was transferred to the agrobacterial strain pGV2260 and transformed B.napus plants produced by agrobacterial transformation essentially as described in Moloney M et al., (1989) Plant Cell Reports 8, 238-242.

Resistance to podshatter was measured using an impact pendulum device (Liu X-Y. 15 Macmillan RH and Burrow RP 1994 Journal of Texture Studies 25 p179-189) (Table 1). The mean energy values shown in Table 1 represent the energy required to rupture the pod on impact with the pendulum. These values are an average from measurements of 20 mature pods. The letters A to L indicate grouping of transformants with significantly different podshatter resistance (ie Group A is 20 significantly different from B when analysed by ANOVA using a Fisher PLSD analysis with a significance level of 95% (Statview 512+). Lines with a number of letters are not significantly different from other lines sharing the same letter. The results shown in Table 1 indicate that 24 lines exhibited significantly higher resistance to podshatter than non-transformed controls whilst 17 lines were not significantly different from the control. The degree and frequency of pod shatter 25 resistance achieved with pWP357-SCV was greater than that obtained by transformation with constructs that downregulate a single DZ-expressed gene.

Table 1

```
PLANTID MEAN ENERGY
 A213-24
            6.752
 A213-53
            5.203
                        8
 A213-34
            3.864
                           C
 A213-21
            3.573
                           C
 A213-4
             3.54
                           C
 A213-61
            3.516
                           Ç
 A214-30
             3.45
                           C
 A214-27
            3.397
                           Ç
 A213-8
            3.277
                           C
 A213-70
            3.271
                           CD
 A214-13
            3.182
                           CD
                                 Ε
 A213-11
            3.182
                           CDEF
· A213-69
            3.005
                              DEF
                                       G
 A213-60
            2.945
                              DEF
                                       GH.
 A214-7
            2.843
                              DEF
                                        G H I
 A213-64
            2.687
                              DEF
                                        G
                                          H
 A214-14
            2.513
                              0 E
                                    F
                                        G H I
 A213-28
            2.581
                              DEF
                                        G H I
 A213-9
            2.547
                                 EF
                                        G
                                          Н
 A213-42
            2.442
                                        G
                                          Н
 A213-31
            2.431
                                        G H
 A214-TO
             2.42
                                          HI
                                                J
 A213-38
            2.295
                                          Н
                                             ŧ
                                                J
 A214-8
             2.26
                                          Н
                                             I
 A213-19
            2.213
                                             ı
                                                J
                                                    K
 A214-25
            2.128
                                             Į
                                                J
                                                    K
 A213-38
            2.059
                                             I
                                                J
                                                    K
 A213-16
            2.005
                                                J
                                                    K
 A213-63
            1.928
                                                J
                                                    K
                                                       L
 A213-33
             1.91
                                                J
                                                    K
                                                       L
 A213-32
             1.901
                                                    K
                                                       L
 A213-37
             1.791
                                                    K
                                                        L
RV27CONT
             1.787
                                                    κ
 A213-10
             1.623
                                                        L
 A213-58
          1.595
                                                        L
 A214-24
             1.55
 A213-3
             1.518
                                                        L
 A213-27
             1.495
 A213-40
             1.395
 A213-47
             1.315
                                                        L
 A213-43
             1.286
                                                        Ĺ
 A213-30
             1.241
```

REFERENCES

- Bleasby A J and Attwood T K, OWL A Non-redundant Composite Protein
 Sequence Database. Nucleic Acid Research 22:3574-3577 (1994)
 - 2. Christoffersen and Laties, Proc. Natl. Acad. Sci. 79, 4060-4063 (1982)
- 10 3. Dean et al., EMBO J. 4: 3055-3061 (1985)
 - 4. Deveraux et al., Nucl. Acids Res. 12:387 (1984)
- Hocking P J and Mason L: Accumulation, distribution and redistribution of
 dry matter and mineral nutrients in fruits of canola (oilseed rape) and the
 effect of nitrogen fertiliser and windrowing. Australian Journal or
 Agriculture Research 44: 1377-1388 (1993)
 - 6. Hakwa et al., Science 198:105-63 (1977)

20

- 7. Jenkins et al: 5th International Conference of Plant Molecular Biology,
 Abstract 310 (1997)
- Johnson-Flanagan AM and Spencer MS: Ethylene production during
 development of mustard (Brassica juncea) and canola (Brassica napus) seed.
 Plant Physiol 106: 601-606 (1994)
 - 9. Liang P and Pardee A: Differential display of eukaryotic messenger RNA by

means of the polymerase chain reaction. Science 257: 967-971 (1992)

- Laing P, Bauer D, Averboukh L, Warthoe P, Rohrwild M, Muller H,
 Strauss M, Pardee A B: Analysis of altered gene expression by differential display. Methods in Enzymol 254: 304-321 (1995)
 - 11. Meakin P J and Roberts J A: Dehiscence of fruit in oilseed rape (Brassica napus L.): anatomy of pod dehiscence. J Expt. Bot 41: 995-1002 (1990a)
- 10 12. Meakin P J and Roberts J A: Dehiscence of fruit in oilseed rape (Brassica napus L.): the role of cell wall degrading enzymes and ethylene. J Expt. Bot 41: 1003-1011 (1990b)
 - 13. Moloney M et al; Plant Cell Reports, 8, 238-242 (1989)

15

5

- 14. Needleman and Wunsch J. Mol. Biol. 48:443 (1970)
- 15. Smith and Waterman Adv. Appl. Math 2: 482 (1981)

CLAIMS

1. Nucleic acid encoding a signal transduction protein involved in the process of dehiscence.

5

- 2. Nucleic acid as claimed in claim 1 wherein the process involves the production of a hydrolytic enzyme.
- Nucleic acid as claimed in claim 1 or claim 2 which is naturally expressed in a dehiscence zone.
 - 4. Nucleic acid encoding a protein wherein the protein:
 - a) comprises the amino acid sequence shown in Figure 1 or;

15

b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above and has at least 40% amino acid sequence identity therewith; or

20

- c) is a fragment of a protein as defined in a) or b) above, which is at least 10 amino acids long.
- Nucleic acid as claimed in any one of claims 1 to 4 which comprises the sequence set out in Figure 1 or a fragment thereof which is at least 30 bases
 long.
 - 6. Nucleic acid, as claimed in any one of claims 1 to 5 in combination with one or more further nucleic acid sequence which is dehiscence-zone expressed.

25

- 7. Nucleic acid which is antisense to nucleic acid as claimed in any one of claims 1 to 6.
- Nucleic acid as claimed in any one of claims 1 to 7 including a promoter or other regulatory sequence which controls expression of the nucleic acid.
 - 9. Nucleic acid which is the naturally occurring promoter or other regulatory sequence which controls expression of nucleic acid as claimed in any one of claims 1 to 8.
 - 10. Nucleic acid as claimed in any one of claims 1 to 9 which is in the form of a vector.
- 15 11. A cell comprising nucleic acid as claimed in any one of claims 1 to 10.
 - 12. A plant cell as claimed in claim 11.
- 13. A process for obtaining a cell as claimed in claim 11 or claim 12 comprising introducing nucleic acid as claimed in any one of claims 1 to 10 into said cell.
 - 14. A plant or a part thereof comprising a cell as claimed in claim 11 or claim 12.
 - 15. Propagating material or a seed comprising a cell as claimed in claim 11 or claim 12.

16. A process for obtaining a plant or plant part as claimed in claim 14 or claim 15 comprising obtaining a cell as claimed in claim 11 and growth thereof or obtaining a plant, plant part, or propagating material as claimed in claim 14 or claim 15 and growth thereof.

5

- 17. A signal transduction protein involved in the process of plant dehiscence.
- 18. A protein which:
- a) comprises the amino acid sequence shown in Figure 1 or;
 - b) has one or more amino acid deletions, insertions or substitutions relative to a protein as defined in a) above, and has at least 40% amino acid sequence identity therewith; or

15

- a fragment of a protein as defined in a) or b) above which is at least
 amino acids long.
- 19. A protein as claimed in claim 17 or claim 18 which is isolated or recombinant.
 - 20. A process for regulating/controlling dehiscence in a plant or a part thereof, the process comprising obtaining a plant or part thereof as claimed in claim 14.

25

21. A process as claimed in claim 20 which comprises obtaining a plant cell as claimed in claim 21 or part of a plant as claimed in claim 14 and deriving a plant therefrom.

- 22. A process as claimed in claim 20 which comprises obtaining propagating material or a seed as claimed in claim 15 and deriving a plant therefrom
- 23. A process as claimed in claim 20 wherein the dehiscence is of a pod or of an anther.
 - 24. Use of nucleic acid as claimed in any one of claims 1 to 10 in the regulation/control of plant dehiscence.
- 10 25. Use of nucleic acid as claimed in any one of claims 1 to 10 as a probe.
 - Use of nucleic acid as claimed in any one of claims 1 to 10 in the production of a cell, tissue, plant part thereof or propagating material.
- Nucleic acid comprising one or more of the underlined sequences as set out in Figure 1, or one or more of the primer sequences in Figure 5, 9 and/or 11.
 - 28. Use of the nucleic acid as claimed in claim 27 as a PCR primer.
- 20 29. Use of a protein as claimed in any one of claims 17 to 19 as a probe.

NUCLEIC ACID AND PREDICTED PROTEIN SEQUENCE OF DZ2

Ncol

-

1/27 120 180 240 300 360 9 CCACCGTGACGGCGCTCATCTTTTGACCTTATCCTAATGGATAAAGAAATGCCCGAGAG GGATGGTGTTTCGACAACTAAGAAGCTAAGAGAAATGGAAGTGAAGTCAATGATTGTTGG GGTGACTTCACTGGCTGACAATGAAGAGGAGCGCAGGGCTTTCATGGAAGCTGGACTTAA GGCACGAGCAGAATCGAAGATGGCAACAAAATCCATGGGAGATATCGAGAAAATAAAGAA GAAACTAAACGTGTTGATCGTCGATGATGATCCACTAAACCTTATAATTCATGAGAAGAT CATCAAAGCGATTGGGGGTATTTCACAGACAGCGAATAACGGTGAGGAGGCAGTAATCAT **[1**] DZ2FL → H K Σ ы ഗ 回 ы × [1] × > Н z Σ [L] 4 ᆸ Σ Σ Н α, ы K ĸ Ωы Ø Ω [x] × K Ω S Гī × > S Z |--| H Þ > Ü S П z K 241 121 181 301 61

420 480 TATGTATGCATAGATACTTGCATGTGTGTTTTTAGAATTTAGGGTTCTTTATCGTCCGT 540 CCATTGCTTGGCAAAACCGTTAACCAAGGACAAGATCATCCCTCTCTATTAACCAACTCAT GGATGCTTGATGGATATATTTTATATTATGGAAACACACATAATAACGTCTAAGTGTG N ← DZ2RL z ᆸ Н × Ω × H Н щ × K Н 361 421

2/27

009

GATATATAATCATGTAGTTGTTGCTTTAAGCTTATAAAATATTAAATAATAAGGGTTTCCT

The primer sites for DZ2FL and DZ2RL are underlined, as are the recognition sequences for Ncol and HindIII restriction enzymes. Shown in bold are the conserved amino acid residues required phosphorylation. The extent of the original PCR product isolated by differential display is shown by \downarrow

FIG. 1 contro

481

CTACC

601

541

WO 99/49046

3/27

Alignment of the predicted protein sequence of DZ2 with those of bacterial response regulator proteins.

```
50
 DZ2
      MATKSMGDIE KIKKKLNVLI VDDDPLNLII HEKIIKAIG. GISQTANNGE
 OMPR
      PHOB
      ...... ....MARRILV VEDEAPIREM VCFVLEQNGF .QPVEAEDYD
NTRC
      ..... .. .. .. .. .. MQRGIVWV VDDDSSIRWV LERALAGAGL .TCTTFENGN
       SPOOF
 CHEY
       ..... MADKELKFLV VDDFSTMRRI VRNLLKELGF NNVEEAEDGV
 ETR
       ..... TKVLV MDENGVSRMV TKGLLVHLGC EVTTVSSNEE
      51
                                                        100
 DZ2
      EAVIIHRDGG SSFDLILMDK EMPERDGVST TKKLREMEVK SM..IVGVTS
 OMPR QMDRLLTR.. ESFHLMVLDL MLPGEDGLSI CRRLRSQS.. NPMPIIMVTA
PHOB SAVNQLNE.. PWPDLILLDW MLPGGSGIQF IKHLKRESMT RDIPVVMLTA
NTRC EVLAALAS.. KTPDVLLSDI RMPGMDGLAL LKQIKQ..RH PMLPVIIMTA
SPOOF
      QALDIVTK.. ERPDLVLLDM KIPGMDGIEI LKRMKV..ID ENIRVIIMTA
CHEY
      DALNKLQA.. GGYGFVISDW NMPNMDGLEL LKTIRADGAM SALPVLMVTA
 ETR
     ....CLRVVS HEHKVVFMDV CMPGVENYQI ALRI..... .HXPLLVALS
      101
                                                        150
 DZ2
      LADNEEERRA FMEAGLNHCL AKPLTKDKII PLINQLMDA. .....
 OMPR KGEEVDRIVG .LEIGADDYI PKPFNPRELL ARIRAVLRRQ ANELPGAPS.
 PHOB RGEEEDRVRG .LETGADDYI TKPFSPKELV ARIKAVMRRI SPM.....
NTRC HSDLDAAVSA .YQQGAFDYL PKPFDIDEAV ALVERAISHY QEQQQPRNIE
SPOOF YGELDMIQES .KELGALTHF AKPFDIDEIR DAVKKYLPLK SN.....
 CHEY EAKKENIIAA .AQAGASGYV VKPFTPATLE EKLNKIFEKL GM.....
 ETR GNTDKSTKEK CMSFGLDGVL LKPVSLDNIR DVLSDLL.......
     151
  DZ2
 OMPR ....
                   OmpR = E. coli (Involved in osmoregulation)
 PHOB
      . . . .
                   PhoB = E. coli (Involved in phosphate utilisation)
                   NtrC = S. typhimurium (Involved in nitrogen utilisation)
 NTRC VNGP
SPOOF
                   SpoOF = B. subtilis (Involved in sporulation)
 CHEY
                   CheY = E. coli (Involved in chemotaxis)
  ETR ....
                   ETR = A. thaliana ETR1 gene encoding an ethylene receptor
                        (partial amino acid sequence)
```

The predicted protein sequence of DZ2 is shown in bold as are the conserved amino acid residues required for phosphorylation of the protein

FIG.2

4/27

Expression analysis of DZ2 in various plant organs using Northerns




FIG. 3

Comparison of bacterial two-component regulatory systems with DZ2

Bacterial two-component system Response regulator Sensor G1 F G2 DD DK N Н Input Histidine protein Receiver Output domain domain domain kinase domain Arabidopsis thaliana ETR1 gene **G1 F G2** DD DK H N Arabidopsis thaliana ERS gene **G1 F G2** DZ2 DD DK

FIG. 4

<u>-1</u>6.5

TATATAAATACGGTTTAACAGATATGTTCTGGTTATAAATGTAATTCNATGTGCCNNTCAANTTTATTTTNATTNGT	78
INTACTAGGGACATTAGTTTTAACNTTTTATATATCATGTAAAAAAAAAA	156
GCAATTATTCTTATAGTGTTTTCTTTTTCCAGAAATTTGACGACAACCTAACTAA	234
AAGTAATTTATATAGATAAATTGAGCAAGCACATTACGAACTGCGGATCAAGGAGAGTCACAATTTAATTCTTA GGC <u>TCTAGA</u> CGAACTGCGGAGCAAGG -> DZ2BGENF XbaI	312
CGTTATACACAAAATTATCTAAATACTATATATATATACAGCTGCATGCTACGATAATGATCAAATGTTTATGTACTT	6 / 27 66 66
TTCAGCGAAAATTCTTGTCGCCATACATTACTGTGTTAATGAATCATTAAATATGTGAAGGAGGAAAAGAGTACAAAA	468
GGAGITTITGTTGAGGCATTTCGCAGACACTGAAATGTGAATAATAATAAAGGAATTGCCGAATTGATTTCTAGTTGGT	546
GAAGTGGGTGAAAATTGTATGTCCATTGCTTATAAACTATAAAATATATAT	624
INNGATAGACCCTAGCTAAAATTTTTAAAATTATACATTCATT	702
TAAAATTGTTTAAGAAACCATTACAAACTCAGCTTGTGGACTCTGAGAGAAACTAAGAGCTAGACATACGGTTAGTAG	780
TGTAGCCGCATTTTTTATGCTTAATTTTGCTTAAGCATGACTTTCTATGCTCCTTTGATGATATTTTTAATATCCTAG	858
GACATATGGATTTGATAAAGATCTTATCAACCATTCAACAAGACCATTAGCTCAACAAACA	936

H Z

I

×

DOTO

1560	NCOI <- GTTCGAAGAAAAAAACCGA <u>CCATGG</u> CAG DZ2BGENR GTGTGAAGATTGTTCGAAGAAAAAAAAAACCGAAGATGGCAACAACGTCAACGTCAACAGGAGATATCAAGAAAACCAAG M A T T S T G D I K K T K
1482	CGCATGCATTAGAACACTTGTATGTTAATTTCCATAATGTTTTGCATAAACATTCTTCGTTTTTAATTAGCTTCTTTTTT
1404	TTATTATTCATAATACAAACACGAAGCCATGCAAGAAGATAGTTACACGCTCATAACAAAAAAAA
1326	AATCTTCTTTAGAATCCAAAATTGTAAGCCGCTTTCTAATCTCTTTTCTAGTATACATATGTAATATGTATAGCATATA
1248	GGTGTGAAAATATGCCTCTTTTCACTTTTTTTCAACTATAAATTTCGATCGA
1170	GTTTGGTAAGACAACTCAGTTGCAACGATGCAGATTACATTTCAGGAAGATTCATGTAAGAAAGA
1092	ATTAATATATTTTTCACTTTTTGTTTTCTTAGGATTTTTTTT
1013	AAICTIGGITTACAGAAITICITIATGCCAAAAATAICATAATATATATAGAATIICGGITTATGATTAAGATGAATTATITTA 1013

O

TCAGTAGAAGTGAAGAAACTTAACGTGTTGATCGTTGATGATGATACAGTAATTCGTAAACTTCACGAGAATATC

H

z

×

ы

TCAGTAGAAGTG -> DZ2BFL

ACTITICGGCAACTAAGAAGCTAAGAGAAATGAAAGTGACGTCTATGATTATTGGGGTGACGACACTGGCTGACAATGAAGAG TCTTTCGACAACTAAGAAGCTAAGAGAAATGGAAGTGAAGTCAATGATTGTTGGGGTGACTTCACTGGCTGACAATGAAGAG AGGAGGCAGTAATCATCCACCGTGACGGCGGCTCATCTTTTGACCTTATCCTAATGGATAAAGAAATGCCCGAGAGGGATGG AGGAGGCAGTGAACATCCACCGCGACGGCAATGCATCTTTCGACCTTATCCTAATGGATAAAGAAATGCCCGAGAGGAGGATGG ••••••••••••••••••••••••••••••••••• GGCACGAGCAGAATCGAAGATGGCAACAAAATCCATGGGAGATATCGAGAAAATAAAGAAGAAGTAAACTAAACGTGTTGATCGTC TCGTCNAT 30 •••••• A) Dz2B cDNA segence (top) Dzz sequence (bottom) 100 260 250

8/27

300

270

GAGCGCAGGGCTTTCATGGAAGCTGGACTTAACCATTGCTTGGCAAAACCGTTAACCAAGGACAAGATCATCCTCTCATTA

380

350

9/27


AACGICATACACCIGIGIG--IGIAIGCATAGATATCTATCCGCAIGIGIGTITTTTAGGGTIGITATGTTIGATTITATIG --AGTGTGTATGTATGCATAGATACTTGCAT--GTGTGTGTTTTTAGAATTTTAGGGTTCTTTATCGTCCGTG --GAAACACACATAAT acaateteatggatgettgatggatggatgaattgtegeeactaataatettatataeattatacattatacaaatatgaaaaaacacatataat 460 ACCAACTCATGGATGCTTGATGGATATATATTTTTATATTATG--••••• 360 440 430

510

500

490


AACGTCTA--



11/27

FIG. 7

NORTHERN ANALYSIS OF EXPRESSION OF DZ2B IN PODS AND OTHER TISSUES



DZ = POD DEHISCIENCE ZONE (20 - 06 daa)

NZ = POD NON-ZONE


AZ = ABSCISSION ZONE

F = FLOWER

L = LEAF

R = ROOT

S = SEED



SUBSTITUTE SHEET (RULE 26)

F16.9

78 ATATATGTGATACAGATACATCTATATACAAATTAAACACGAAACCATACATGCACGGTGTGATCACACACGCACACA 234 TCTTGAGAGAAAAGAAATCGAAGATGGCAACAAAATCCACCGGAGGTACCGAGAAAACCAAGTCGATAGAAGTGAAAGA × ۲ × 口 <u>[</u>-Ö Ç × EH 4 Σ

13/27

312 AGAAACTAATCAACGTGTTGATCGTCGATGATGATCCATTAAACCGTAGACTCCACGAGATGATCATCAAAACGATCG <- CGATCG Σ ш ;;; П K Z Д Ω П >

<- CGAATAACGGTGAGGAGGCATTAATCA GW1</p>


390 GAGGAATTTCTCAGACTGCAAAGAATGGCGAAGAGGCNGTGATCCTCCACCGTGACGGCGAAGCATCTTTCGACCTTA GAGGAATTTCTCAGACTGCAA AT3GW2

ល ⋖ O Ω ፈ 二 Н > O Z × A, O S O

SUBSTITUTE SHEET (RULE 26)

TTCTAATGGATAAGGAATGCGTGGGATTTCGACAATTAAGANGCTAAGAGAAATGAAAGGGACGTCAA L M D K E M P E R D G V S T I K X L R E M K G T S M	468
TGATCGTTGGGGTAACGTAGCTGACCAAGAAGAGCGTAAGGCTTTTATGGAAGCTGGGCTCAACCATTGCT I V G V T S V A D Q E E E R K A F M E A G L N H C L	546
TGGAAAAACCCTTAACCAAGATCTTCCCGCTCATTAGCCACCTCTTCGATGCTTGATGGATG	14/27 74/27
AATGTATCTATATTTTCAATCATGAAATCACCTACACGTGTATTTGACACAAAAATCTGCATTTGTTGTGATATAGGG	702
TITCTCATATCTATGTTTGATTTTTTTTTTTTCTTCGTCCGAGGTAAAATCATGCAAGTCATTTCTTTTGGCTAATAAAA	780

FIG. 9 contro



16 / 27




FIG. 10 contro

LSATKKLREMKVTSMI IGVTTLADNEEERKAFMEAGLNHCLAKPLSKAKILPLINNLMDA

80

9

DZ2AT3 peptide sequence (top) DZ2B peptide sequence (bottom)

a

FIG. 11

AAAACAGTTTCACGATTGTTTTTTTTTTTTTTTTTAATTTTTAACATGTATGGGGATATTTGGCTATAAATTGGCTATAAAAA TAAAAAAATTTCACGATAGATTGTTGAATTTTTTTTTT
234 312 390 468 546 624
312 390 468 546 624
390 468 546 624

Ø

出

U

回

耳

Н

ĸ

pc,

Z

H

Д

Ω

Ω

Ω

>

Z

L

×

×

H

Ö

Ö

E


ß

×

E


Σ

TGAAGAAGAAACTAATCAACGTGTTGATCGTCGATGATGATCCATTAAAACCGTAGACTCCACGAGTGTCATCAAAA




SUBSTITUTE SHEET (RULE 26)

20/27



SUBSTITUTE SHEET (RULE 26)

21/27 Sphl IllbniH Itsq **Σ∞H**Λ IUds CaMV polyA ЮЧХ RB Over/ drive Clai EcoRI Sstl D22 looM pDZ28-EcoRI EcoRI Nos polyA Sphl RK2 ori BamHi Xpsl **Hms8** Smal Gm/Km Kpnl lR∞∃ Sstl IUds nptII Kpnl Itsq Sstl 1Hms8 Itsq pDZ2B-DZ2as-SCV pNos ΛΙ∞3 ApB Sphl ColE1 ori Clai



23 / 27

240 300 360 $\begin{array}{c} 480 \\ 112 \end{array}$ 540 132 $600 \\ 152$ GGCATCACGAGGGTACCCGTAAATCCCACCATACAACAAAGTTCTGTGAAAGTCTCCCAA GGTCATGAAGATGGAAGCTTCGAAACCGATAGTTTAATCAAGCTCAACAACGACGACGAC GTTCTTACCTTGAAAAGCTCCGATAGACCCACTACCGAATCATCAACTGTTAGTGTTTCG AACTTCGGAGCAAAAGGTGAAAAACCGATGATACTCAGGCTTTCAAGAAAGCATGG N F G A K G D G K T D D T Q A F K K A W A A A C T G C A A A G A G T C C A T A T T A G T T C T T T C T C A G A A A A A A A A C A G T G T T T C T G A A A A G A TT A G C G TT T C A A A C C C C G A A A T G G C C C G T T G T A G G T C T T G C T A T T T T T T T T A TGCGTTCTTTTGATGCTCGCTTGCTGCCAAGCTTTGAGTAGCAACGTAGATGATGTAT
C V L L M L A C C Q A L S S N V D D G Y **AAGAAGGCATGTTCAACAAATGGAGTGACTACTTTCTTGATTCCTAAAGGGAAGACTTAT** GGCACTTTATCAGCTTCTACAAACGATCGGATTACAGTAATGACAAGAACCACTGGCTT G T L S A S T K R S D Y S N D K N H W L CTCCTTAAGTCTATTAGATTCAGAGGCCCATGCAAATCATTACGTAGCTTCCAGATCCTA S S O S s ပ П S R P ල G S 24 S

1 3 XX

1080 720 780 840 232 006 252 ATTITGGAGGACGTTAATAATCTATCAATCGATGGCGGCCTCGGCGGGGATTGTTGATGGC GCGCCAACGGCTCTTACTCTACAACCTAAACAATTTGAATGTGAAGAATCTGAGAGTG A A A A A C A T T C G A A T C C C A A T T C A G A C A T T G G G A C A G G T G A T T G C A T T C C A T T G A G GATGGATCGCAAAATGTTCAAATCAATGATTTAACTTGCGGCCCCGGTCATGGCATCAGC D G S Q N V Q I N D L T C G P G H G I S AGAAATGCACAGCAGATTCAGATTTCGATTGAGAAATGCAACAGTGTTGATGTTAAGAAT GTTAAGATCACTGCTCCTGGCGATAGTCCCAACACGGATGGTATTCATATCGTTGCTACT **ATTGGAAGCTTGGGGATGACAATTCCAAAGCTTATGTATCGGGAATTAATGTGGATGGT** ပ Ξ S Ω Z ပ G H G × Z H G Z Z Ś S S S Z \circ G 0 z **-**--0 S S

24/27

A

G G S G 332	ATCCGATCATA 1200 N P I I 352	CGGTTCAAGTG 1260 A V Q V 372	CGATAATGTTT 1320 A I M F 392	ACATCAAAGGA 1380 N I K G 412	CTCCTAAATGC 1440 S P K C 432	rggttatagat 1500 433	FAAGGATTTAT 1560	ACTGAAGTTGA 1620	1657
GCTACGCTCTCTGAGACTGAGAGTAAGAATCAAGACTTACCAGGGAGGG	ACTGCTAAGAACATTCCAAAACATTCGTATGGATAATGTCAAGAATCCGATCATA T A K N I K F Q N I R M D N V K N P I I	ATCGACCAGAACTGCGACAAGGACAAATGCGAACAACAAGAATCTGCGGTTCAAGTG I D O N Y C D K D K C E Q Q E S A V Q V	AACAATGTCGTCTATCGGAACATACAAGGTACGAGCGCAACGGATGTGTTT N N V V Y R N I Q G T S A T D V A I M F	AATTGCAGTGTGAAATATCCATGCCAAGGTATTGTGCTTGAGAATGTGAACATCAAAGGA N C S V K Y P C Q G I V L E N V N I K G	GGAAAAGCTTCTTGCAAAATGTCAATGTTAAGGATAAAGGCACCGTTTCTCCTAAATGC 1440 G K A S C K N V N V K D K G T V S P K C 432	CCTTAATTACTAAGTTGATTATGTAATATACATAAATACGTATTATATGTGGTTATAGAT P	ATCTACGTATTGATTCTCGATATATATAGAAAACTAAGGATTTAT	ATAGTTGAGATAATTGTTGTTGTATATGGTTCACTGAAGTTGA 1620	retcatitete FIG. 15conto
GAGACTGACAATGGAGTA E T D N G V	ATTAAATTCCAAAACATT I K F Q N I	TACTGCGACAAGGACAAA Y C D K D K	ratcggaacatacaaggt _/ Y R N I Q G	AAATATCCATGCCAAGGTA K Y P C Q G	FGCAAAATGTCAATGTT/ C K N V N V	AGTTGATTATGTAATATA(ttgcttgtccacgaataatgaataatgtcatttgtc FIG. 15con
GCTACGCTCTCT A T L S	ACTGCTAAGAAC T A K N	ATCGACCAGAA(I D O N	AACAATGTCGTC N N V V	AATTGCAGTGTG N C S V	GGAAAAGCTTCT' G K A S	CCTTAATTACTA P	GCCATCTATATCCTT	GGGAATATACATACA	TTGCTTGTCCAC

aggtgaccgttgctgatggcaatgtgctggtcaagcgagaggtagacggtggcttggagaNVLVK R E V D G cagttaaagtcaaattgccagctgtcattagcgccgacttgcggctcaatgagccgcggt K V K L P A V I S A D L R L N E P R acgctactctgcccaatatcatgaaggccaagaagaagcccatcaaaaagctcacagcca ATLPNIMKAKKKPIKKLTAT cagatgtcggtgtggacttggcgccacgtcaacaagtgttgagcgtagaagacccgccca G V D L R Q QE Q A G S I V P D V D T L Ι T K L agggtcatttgtaatgcaatgtcaccaatacagttgttttagttcttacaaattcttcgt G H L * gaggttttcagctgttaccaataatattttttcaaaaatcgattttattttacttgtaatt taaaagatcaaatattaatacaatgaacatttttgtaacagcaatctttttttatattt

atataataaagtatgttttctggtcaaaa

FIG. 16

tatactctccata Y T L H T • &> S gttgggtcgaaccataggtggaaagcttctttttttctctcgcttgacaaatcctctggt L G R L L S L S L D R S S Gctt aca atggttcg M V R Ø gaaa cgggaac G T aat cacgaaggctctggagacaaagaacagcagtttcatct ttca gaacagtcacaacattctatcttaaatcac TVTTFYLKSP atcgatttcgagttcttgggaaacataagtggccatcct I D F E F L G N I S G H P Y a 8 8 4 agctge A E catggtaaa (G K ctcta L Y cgggttttcagtcccatcaggagttt tccctggtaactctgctg Ø [→ Ú tttac Y age ctaatgt N V gatga D E 881 00 5

-16. 17

3

ccgacc D

306bp

0SR79

Appl. No. 09/646,679 (U.S. Nat'l Phase of PCT/GB99/00905) Docket Number: 0623.0890000 K&S Ref: KVC/PS/CF/P20895US

POWER OF ATTORNEY FROM ASSIGNEE

The Assignee hereby appoints the following U.S. attorneys to prosecute this application and any continuation, divisional, continuation-in-part, or reissue application thereof, and to transact all business in the U.S. Patent and Trademark Office connected therewith: Robert Greene Sterne, Esq., Reg. No. 28,912; Edward J. Kessler, Esq., Reg. No. 25,688; Jorge A. Goldstein, Esq., Reg. No. 29,021; Samuel L. Fox, Esq., Reg. No. 30,353; David K.S. Cornwell, Esq., Reg. No. 31,944; Robert W. Esmond, Esq., Reg. No. 32,893; Tracy-Gene G. Durkin, Esq., Reg. No. 32,831; Michael A. Cimbala, Esq., Reg. No. 33,851; Michael B. Ray, Esq., Reg. No. 33,997; Robert E. Sokohl, Esq., Reg. No. 36,013; Eric K. Steffe, Esq., Reg. No. 36,688, Michael Q. Lee, Esq., Reg. No. 35,239; Steven R. Ludwig, Esq., Reg. No. 36,203; Raz E. Fleshner, Esq., Reg. No. 34,331; John M. Covert, Esq., Reg. No. 38,759; and Linda E. Alcorn, Esq., Reg. No. 39,588. The Assignee hereby grants said attorneys the power to insert on this Power of Attorney any further identification that may be necessary or desirable in order to comply with the rules of the U.S. Patent and Trademark Office.

Send correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C. 1100 New York Avenue, N.W. Suite 600 Washington, D.C. 20005-3934 U.S.A.

Direct phone calls to 202-371-2600.

FOR: Biogemma UK Limited

SIGNATURE:

BY:


Tina Lorraine Barsky

TITLE:

Company Secretary

DATE:

18 December 2000



Declaration for Patent Application

Appl. No. 09/646,679 (U.S. Nat'l Phase of PCT/GB99/00905) Docket Number: 0623.0890000 K&S Ref: KVC/PS/CF/P20895US

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled Signal Transduction Protein Involved in Plant Dehiscence, the specification of which is attached hereto unless the following box is checked:

as United States No. 09/646.679		ernational Application Number <u>PCT</u>	<u>/GB99/00905</u>	(U.S. Appl.
I hereby state that I have		tents of the above identified specific	cation, includ	ing the
I acknowledge the duty t	to disclose information that is mat	erial to patentability as defined in 3	7 C.F.R. § 1.5	i6.
or inventor's certificate, than the United States lis	or § 365(a) of any PCT internation sted below, and have also identified	119(a)-(d) or § 365(b) of any foreignal application, which designated at ed below any foreign application for g date before that of the application	t least one cou r patent or inv	ntry other entor's
Prior Foreign Application Claimed	n(s)			Priority
9806113.8 (Application No.)	Great Britain (Country)	20 March 1998 (Day/Month/Year File	⊠ Yes ed)	□ No
(Application No.)	(Country)	(Day/Month/Year File	□ Yes ed)	□ No
I hereby claim the benef	it under 35 U.S.C. § 119(e) of any	y United States provisional applicati	ion(s) listed be	elow.
(Application No.)	(Filing Date)			
(Application No.)	(Filing Date)			
international application claims of this application provided by the first par patentability as defined	designating the United States, lis is not disclosed in the prior Unit graph of 35 U.S.C. § 112, I ackn	United States application(s), or under sted below and, insofar as the subjected States or PCT international appl nowledge the duty to disclose informational between the filing date of the	at matter of ead ication in the s mation that is r	ch of the manner naterial to
PCT/GB99/00905 (Application No.)	March 22, 1999 (Filing Date)	Abandoned (Status - pat	ented, pending	g, abandoned)
(Application No.)	(Filing Date)) (Status - pat	ented, pending	g, abandoned)

Appl. No. 09/646,679 (U.S. Nat'l Phase of PCT/GB99/00905) Docket Number: 0623.0890000 K&S Ref: KVC/PS/CF/P20895US

Send Correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C. 1100 New York Avenue, N.W. Suite 600 Washington, D.C. 20005-3934

Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or

any patent issued thereon. Full name of sole or first inventor Wyatt Paul Date Signature of sole or first inventor Cambridge, Great Britain Citizenship Great Britain Post Office Address Biogemma UK Limited, 200 Science Park, Milton Road, Cambridge, CB4 OGZ, Great Britain Full name of second inventor Jeremy Alan Roberts 8/12/00. Date Signature of second inventor Residence Leicestershire, Great Britain Citizenship Great Britain Biosciences Post Office Address University of Nottingham, Division of Plant Science, School of Biological Sciences, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, Great Britain Full name of third inventor Catherine Whitelaw Date Signature of third inventor Residence Beltsville, Maryland Citizenship Great Britain USDA/ARS/SARL, Room 208, Building 006 BARC-West, 10300 Baltimore Avenue, Beltsville, MD 20705-2350, USA

P:\USERS\PChoi\0623\089\declaration wpd

(Supply similar information and signature for subsequent joint inventors, if any)

The first from the first first

Declaration for Patent Application

Appl. No. 09/646,679 (U.S. Nat'l Phase of PCT/GB99/00905) Docket Number: 0623.0890000 K&S Ref: KVC/PS/CF/P20895US

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled <u>Signal Transduction Protein Involved in Plant Dehiscence</u>, the specification of which is attached hereto unless the following box is checked:

as United State No. 09/646,67	9); and	nternational Application Number]	PCT/GB99/00905	(U.S. Appl.
was amended	on(if app	olicable).		
	e reviewed and understand the coany amendment referred to above	ontents of the above identified spee.	eification, includi	ing the
I acknowledge the duty	to disclose information that is m	aterial to patentability as defined	in 37 C.F.R. § 1.5	66.
or inventor's certificate than the United States I	or § 365(a) of any PCT internat isted below, and have also identi	§ 119(a)-(d) or § 365(b) of any foinnal application, which designate fied below any foreign application and date before that of the application	ed at least one country of for patent or inv	ntry other entor's
Prior Foreign Applicati Claimed	on(s)			Priority
9806113.8 (Application No.)	Great Britain (Country)	20 March 1998 (Day/Month/Year	⊠ Yes Filed)	□ No
(Application No.)	(Country)	(Day/Month/Year	□ Yes Filed)	□ No
I hereby claim the bene	fit under 35 U.S.C. § 119(e) of a	ny United States provisional appli	ication(s) listed be	elow.
(Application No.)	(Filing Date)			
(Application No.)	(Filing Date)			
international applicatio claims of this applicatio provided by the first pa patentability as defined	n designating the United States, I on is not disclosed in the prior Un ragraph of 35 U.S.C. § 112, I acl	United States application(s), or unlisted below and, insofar as the sulnited States or PCT international aknowledge the duty to disclose infavailable between the filing date ion.	bject matter of each application in the reformation that is n	ch of the manner naterial to
PCT/GB99/00905	March 22, 1999	Abandoned		
(Application No.)	(Filing Dat		patented, pending	g, abandoned)
(Application No.)	(Filing Dat	re) (Status -	patented, pending	g, abandoned)

Appl. No. 09/646,679 (U.S. Nat'l Phase of PCT/GB99/00905) Docket Number: 0623.0890000 K&S Ref: KVC/PS/CF/P20895US

Send Correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C. 1100 New York Avenue, N.W. Suite 600 Washington, D.C. 20005-3934

Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

my patent issued mercon.	
Full name of sole or first inventor Wyatt Paul	
Signature of sole or first inventor	Date
Residence Cambridge, Great Britain	
Citizenship Great Britain	
Post Office Address Biogemma UK Limited, 200 Science Park, Milton Road, Cambridge, CB4 OGZ, Great Britain	
Full name of second inventor Jeremy Alan Roberts	
Signature of second inventor	Date
Residence Leicestershire, Great Britain	
Citizenship Great Britain	
Post Office Address University of Nottingham, Division of Plant Science, School of Biological Sciences, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, Great Britain	
Full name of third inventor Catherine Whitelaw	
Signature of third inventor Amitifan , 11/12/00	Date
Residence Beltsville, Maryland	
Citizenship Great Britain	
Post Office Address USDA/ARS/SARL, Room 208, Building 006 BARC-West, 10300 Baltimore Avenue, Beltsville, MD 20705-2350, USA	

P:\USERS\PChoi\0623\089\declaration wpd

(Supply similar information and signature for subsequent joint inventors, if any)

Declaration for Patent Application

Appl. No. 09/646,679 (U.S. Nat'l Phase of PCT/GB99/00905) Docket Number: 0623.0890000 K&S Ref: KVC/PS/CF/P20895US

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter that is claimed and for which a patent is sought on the invention entitled Signal Transduction Protein Involved in Plant Dehiscence, the specification of which is attached hereto unless the following box is checked:

Ø	as United States No. 09/646,679)		national Application Number PCT/GF	399/00905 (U.S. Appl.
		reviewed and understand the contry amendment referred to above.	ents of the above identified specification	on, including the
I ackno	wledge the duty to	disclose information that is mate	rial to patentability as defined in 37 C.	F.R. § 1.56.
or inver	ntor's certificate, on turited States list ate, or PCT interna	r § 365(a) of any PCT internation and below, and have also identified	119(a)-(d) or § 365(b) of any foreign a al application, which designated at lead below any foreign application for pat date before that of the application on v	st one country other ent or inventor's
Prior Fo	oreign Application d	n(s)		Priority
980611 (Applie	3.8 ation No.)	Great Britain (Country)	20 March 1998 (Day/Month/Year Filed)	⊠ Yes □ No
(Applic	ation No.)	(Country)	(Day/Month/Year Filed)	□ Yes □ No
I hereby	y claim the benefit	t under 35 U.S.C. § 119(e) of any	United States provisional application(s	s) listed below.
(Applic	cation No.)	(Filing Date)		
(Applie	ation No.)	(Filing Date)		
internate claims of provide patental	tional application of this application and by the first para bility as defined in	designating the United States, liste is not disclosed in the prior Unite graph of 35 U.S.C. § 112, I ackno	nited States application(s), or under § 3 and below and, insofar as the subject may ded States or PCT international applicational application of the duty to disclose informational allable between the filing date of the p.	on that is material to
	B99/00905	March 22, 1999 (Filing Date)	Abandoned (Status natural)	d, pending, abandoned)
(Аррис	cation No.)	(Fling Date)	(Status - patente	u, penumg, avandoned)
(Applie	cation No.)	(Filing Date)	(Status - patente	d, pending, abandoned)

Send Correspondence to:

STERNE, KESSLER, GOLDSTEIN & FOX P.L.L.C.
1100 New York Avenue, N.W.
Suite 600
Washington, D.C. 20005-3934

Direct Telephone Calls to:

(202) 371-2600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Wyatt Paul	
Signature of sole or first inventor W. Paul. 77/12/09.	Date
Residence Cambridge, Great Britain	
Citizenship Great Britain	
Post Office Address Biogemma UK Limited, 200 Science Park, Milton Road, Cambridge, CB4 OGZ, Great Britain () BW	
Full name of second inventor Jeremy Alan Roberts	
Signature of second inventor	Date
Residence Leicestershire, Great Britain	
Citizenship Great Britain	
Post Office Address University of Nottingham, Division of Plant Science, School of Biological Sciences, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, Great Britain	
Full name of third inventor Catherine Whitelaw	
Signature of third inventor	Date
Residence Beltsville, Maryland	
Citizenship Great Britain	
Post Office Address USDA/ARS/SARL, Room 208, Building 006 BARC-West, 10300 Baltimore Avenue, Beltsville, MD 20705-2350, USA	

P:\USERS\PChor\0623\089\declaration.wpd

(Supply similar information and signature for subsequent joint inventors, if any)

The first and the first first and the first

100